

Trachoma Matters



8000BC

2020

# Trachoma


Through History

Special Edition: by Katherine Schlosser



international  
**trachoma**  
initiative

Dedicated to eliminating  
the world's leading cause  
of preventable blindness



Surgery preparation

**Trachoma** is a **preventable** disease that affects the eye and can lead

to blindness if left untreated. Repeated infections scar and distort the inner upper eyelid, causing the eyelashes to turn inwards and scratch the cornea, ultimately leading to blindness.

Trachoma has been causing blindness for several millennia. As scientists have only gained a thorough understanding of the disease within the last several decades, the history of trachoma cannot be pinpointed precisely. However, the history of ophthalmias, eye diseases with visible symptoms, is frequently documented. Through these records and a modern understanding of trachoma, the history of this ancient disease can be constructed.

The struggle against trachoma began as early as 8000 B.C. Over these thousands of years, trachoma has blinded great thinkers, traveled across continents, and inspired a vigilant medical inspection process. Today, trachoma continues to blind millions in the developing world, but great progress has been made towards elimination. In the past 50 years, scientists cultured the agent, discovered antibiotics to treat active infection, and developed the **SAFE** strategy, which combines treatment and prevention. Although packaged as a comprehensive approach to trachoma control in the past several decades, the SAFE strategy is truly the culmination of a struggle against trachoma that began thousands of years ago.

## Trachoma and the **Ancient** World

The most probable geographic origins of trachoma can be established by analyzing the prevalence rates among different racial groups. After looking at the distribution of trachoma among different races, researcher Taborisky determined that Central Asia is most likely the place where trachoma originated.<sup>55, 63</sup> In the first half of the twentieth century, when Taborisky was conducting his research, people of Central Asian ancestry had a higher rate of trachoma than people whose origins lay elsewhere. The main tribes from Central Asia were the Mongols and the Finns. Groups of people who came into contact with the Mongols and the Finns had rates of trachoma correlating to the amount of contact. As Taborisky explains, “nearly always people who contracted the disease suffered from it less than those who propagated it.”<sup>55</sup> Taborisky was not alone in his assessment. Arthur MacCallan, who also conducted research on trachoma, agreed that the origins of trachoma were in Central Asia. Perhaps trachoma did not only emerge in Central Asia, as diseases often emerge simultaneously in several areas, but the evidence suggests that Central Asia was one of the regions of origin.<sup>39, 55</sup>

8000 BC

First evidence of trachoma found in Australia

Evidence in China of first written trachoma therapies

2600 BC

1550 BC

The Ebers papyrus in Egypt contains references to trachoma

Trachoma spreads during the Peloponnesian War

431- 404 BC

**Trachoma and the Ancient World**

Although trachoma appears to have originated in Central Asia, one of the oldest possible signs of trachoma has been found in skeletons in Australia. Archaeologists discovered a lesion, an abnormal change in structure due to injury or disease, in the skulls of Australian skeletons, the cause of which had not been identified. The lesions were found frequently, which suggested the cause was common among Australians. The size and depth of the lesion varied slightly among skulls and based on these observations, scientists concluded that the lesion was caused by an infection of the tissue, which penetrated into the bone and not the reverse. Based on the lesion's proximity to different tissues, several eye infections were identified as possible causes. The scientists analyzed the presence of different diseases in Australia's history, as well as the geographic distribution of skulls containing the lesion. The lesion was found in the same areas of Australia where trachoma afflicts Australians today. After completing these analyses, the scientists concluded that trachoma was the most compatible disease. The skulls investigated date back to 8000 B.C. and, if the analysis is correct, it indicates trachoma's presence in Australia at that time.<sup>67</sup>

In addition to the conclusions drawn from these modern observations, there are many written records of trachoma dating back several thousand years. Written therapies for trachoma from 2600 B.C. have been found in China.<sup>28</sup> Another reference is contained in the Ebers papyrus from the 16th century B.C., which is a collection of medical prescriptions found in Egypt.<sup>63</sup> Trachoma was also present in Ancient Greece and Rome. Several written records of the disease have been found from this era (1200 B.C. to 400 A.D.). The Hippocratic Corpus, the collection of work from Hippocrates' school of medicine (5th century B.C.), contains sections about trachoma. Several doctors, including Celsus (1st century A.D.), Discorides (40-91 A.D.), Galen (129-216 A.D.), Grassus (12th century A.D.) and Chauliac (1300-1368 A.D.), mention trachoma in their works.<sup>2, 7</sup> Discorides was the first person to use the word "trachoma" around 60 A.D. in his work *Materia Medica*.<sup>4</sup> In addition to references in medical texts, trachoma was also part of popular discourse. Several of the plays by Aristophanes, including *Plutus*, contain allusions to trachoma. In this particular play, Plutus, god of riches, has been blinded by Zeus. This blindness is believed to be a reference to the widespread trachoma in Greece at the time.<sup>63</sup> Scholars also believe the biblical passage about Leban's eldest daughter, which states "Leah was tender eyed,"<sup>58</sup> is describing trachoma.<sup>63</sup> Through these texts and others, the history of trachoma can be traced back thousands of years.

By reading these documents, historians have discovered not only the long history of the disease, but also the extensive record of treatments. Around 1550 B.C. the Ebers papyrus suggested that in order to treat trichiasis, the turning inwards of the eyelashes and scratching of the cornea, one should "pull hairs and apply mixture of myrrh, lizard's blood, and bats blood until healed or a mixture of fly's dirt, red ochre and urine or a mixture of frankincense, lizard dung, ox blood, ass blood, pig blood, hound blood, goat blood, stibium and malachite."<sup>2</sup> Other treatments included scraping of the eyelid with various tools and surgeries to secure the eyelashes away from the eyeball. The principle of removing or curing the infected tissue and of physically fixing the eyelashes away from the eyeball to prevent scarring are a common theme of trachoma treatment.

**Did you know**

**St. Paul, Cicero, Horace and Galileo all had trachoma?**

**200 BC**

Hippocratic Corpus mentions trachoma

Discorides first uses the word trachoma

**60****1095**

Beginning of Crusades accelerates spread of trachoma

## Trachoma, Soldiers and Civilians


AS treatments have changed over time, so too has the prevalence and distribution of trachoma. When civilizations were isolated from one another without extensive interaction, trachoma affected segments of society, but the extent of disease was limited. When mass movements of people occurred, however, trachoma spread beyond the isolated populations it affected. Prior to the 19th century, these mass movements were mainly the result of war. Both the Peloponnesian War (431-404 B.C.) and the Crusades (beginning 1095 A.D.) contributed to the spread of trachoma, and Napoleon's campaigns across Europe and Africa caused trachoma's prevalence to increase rapidly.<sup>63</sup>

When Napoleon began his quest into Egypt in 1798, he not only had to fight the Egyptian army, but also blinding eye diseases that would incapacitate thousands of his soldiers. Military or Egyptian ophthalmia, as the disease was called, infected three thousand of Napoleon's troops in two and a half months, blinding many.<sup>63</sup> Although military ophthalmia was viewed as a single disease, the affliction was actually a combination of several eye infections, including Koch-Weeks conjunctivitis (known today as *Haemophilus influenzae* biotype III), gonococcal conjunctivitis, and trachoma.<sup>2</sup> The combination of these infections often aggravated the symptoms of each and most likely the gonococcal infections were responsible for the rapid onset of blindness.<sup>29</sup>

When Napoleon's troops withdrew from Egypt they spread the ophthalmia, including trachoma, from Egypt, where the disease was endemic, across Europe, where trachoma was only present sporadically. The British soldiers were also infected with trachoma as a result of their military campaigns in Egypt. The British troops spread trachoma on their way home and trachoma became a public health problem in the United Kingdom after their return.<sup>11</sup> Contact between Europe and Egypt through the Napoleonic Wars facilitated the rapid spread of trachoma.

**Napoleon's** campaigns across Europe and Africa caused trachoma's prevalence to increase rapidly.

Napoleon's campaigns, however, did not introduce trachoma into Europe. Trachoma had been present in Europe between 1200 A.D. and 1700 A.D. but had remained largely unnoticed. Military activities allowed trachoma to expand quickly because many people from different areas were coming into contact with one another. The presence of trachoma in Europe at the end of the 18th century is indicated by the troubles faced by other armies. Infected soldiers were recruited into the ranks of the Italian, Russian and Belgian armies, and the infection spread within the cramped military quarters.<sup>11</sup> If infected soldiers were allowed to remain in the army, they spread the infection to their fellow soldiers, yet if these soldiers were discharged, they brought the infection home with them and began civilian epidemics.



Sir Ralph Abercromby's troops return from Egypt, prompting epidemic in England

John Vetch makes sustained case of contagiousness

1798

Napoleon begins his quest into Egypt

1801

1803

Trachoma epidemic begins in Paris

1807

*Gentlemen's Magazine* publishes article about troops faking trachoma

### Trachoma, Soldiers and Civilians


Confusion over how trachoma was transmitted and lack of effective treatment also allowed trachoma to spread rapidly. The concept of contagiousness and the idea of each disease having a unique cause were still topics being debated by doctors in the 19th century. Various environmental factors were often blamed for disease and the idea of person-to-person transmission of disease was not widely accepted.<sup>66</sup> A common belief was that only soldiers who served in Egypt could be afflicted with trachoma. However, when civilians and soldiers who had never been to Egypt began contracting the infection, the contagious nature of trachoma became clear. In 1810, England appointed a team of professionals to investigate trachoma. The team recommended that infected people should be isolated and emphasized cleanliness, including frequent washing of persons and clothing and not sharing towels or linens. These new guidelines resulted in an improvement in the standard of living for many British soldiers.<sup>11</sup> Treatments also included removal of granulations, various eyewashes, and bloodletting, which involved making cuts near the eye and allowing the blood to flow from these cuts until the patient fainted.

While physicians were largely unsuccessful at treating trachoma, they did understand how to prevent the spread of infection. The ophthalmic schools created in England are an example of successful trachoma control. Trachoma was a serious problem in the poorer schools in the late nineteenth and early twentieth centuries in London.<sup>11</sup> The British government created special schools, including Hanwell Ophthalmic School and Swanley, where infected children were isolated.<sup>11, 36</sup> Strict sanitary measures were enforced according to the expert committee's guidelines. The schools had the joint success of stopping the transmission from infected to healthy children and helping the infected children recover from infection without progressing to blindness. The fact that the nurses and staff at the schools did not become infected also helped prove that trachoma was not airborne. Recognizing sanitation as a key to trachoma control helped eliminate trachoma as a problem in England.

Trachoma's impact on society reached beyond the physician's office. When trachoma struck France in 1803, the epidemic began in Paris and spread from there. The prevalence of trachoma was extremely high in Paris and all social classes were affected. Trachoma was labeled a fashionable disease because it was so widespread.<sup>63</sup> In Britain, soldiers attempted to fake ophthalmia to avoid military service. Knowledge of this ploy was widespread and *Gentlemen's Magazine* published an article in 1807 about troops from the 28th regiment of foot soldiers using an ointment to fake trachoma in an attempt to be discharged. The soldiers' plot was exposed when surgeons were sent to investigate by authorities concerned at the high rate of ophthalmia in this region.<sup>63</sup>

The increase in prevalence of trachoma that accompanied the Napoleonic Wars created an atmosphere in which trachoma could no longer be ignored. These epidemics raised awareness and brought money and scientific expertise to the field of ophthalmology. In Britain these epidemics influenced the establishment in 1805 of the Central London Ophthalmic Hospital, which is still an active ophthalmological hospital (Moorfields Eye Hospital) today.<sup>4, 21</sup> In 1903, Sir Ernest Cassel, who financed the Aswan Dam, donated £40,000 to train Egyptian ophthalmologists. Sir Cassel hoped free ophthalmology services would become available to those in need.<sup>39</sup> The Egyptian camel drivers during World War I benefited from Sir Cassel's donation. The high prevalence of trachoma spurred these developments in the field of ophthalmology.

**When Lewis and Clark were traveling on the Columbia River in the United States, they encountered native tribes suffering from ophthalmia, which may have been trachoma.**



## Trachoma and Immigration

At the end of the 19th century and beginning of the 20th century, many immigrants left their homes in Europe and set out for a new life in America. Immigrants infected with trachoma carried the disease with them on their long journey and were a source of infection for the people they encountered. Anxiety about diseases transported by immigrants led to legislation being passed in 1891 by the U.S. Congress that stated, “[t]hat the following classes of aliens shall be excluded from admission into the United States [...] persons suffering from a loathsome or a dangerous contagious disease.”<sup>12</sup> The concern about trachoma was so great that in 1897, trachoma was the first disease classified as a loathsome or dangerous contagious disease by the U.S. government.<sup>31</sup> From 1898 until 1905, immigrants exhibiting symptoms were examined for trachoma, and after 1905, all immigrants wishing to enter the United States were examined.<sup>30, 41</sup> Trachoma was viewed as a serious threat by many governments and this threat guided decisions regarding immigration legislation. As a result of the strict American laws, steamship companies decided to conduct trachoma inspections in Europe before embarkation. Companies refused to transport infected passengers because of the expense of transporting rejected immigrants back to Europe. Many immigrants traveled to the U.S.A. through Canada, the United Kingdom, Germany and other countries. Immigrants who were suffering from trachoma were refused passage to the U.S.A. and remained in port cities where they created trachoma outbreaks. These outbreaks caused several countries, including Canada who passed legislation in 1902, to exclude immigrants with trachoma.<sup>11</sup>



An immigration officer at Ellis Island checks a woman's eyelids for signs of infection.

If an immigrant did pass the preliminary inspections in Europe, they still faced the scrutiny of the United States Public Health Service (USPHS) physicians at Ellis Island or other similar facilities across the country. The physicians had approximately six seconds per immigrant to determine if he or she had any physical or mental ailments which should bar entry to America. The exam included a physician examining the underside of the immigrants' eyelids using either his fingers or a buttonhook to evert the eyelids, a process which was frightening for the immigrants. If an immigrant failed the eye exam, he was sent to a special committee that decided whether or not he could enter the country. Occasionally, if the board thought the immigrant could be cured and had sufficient financial backing to pay for medical treatment, the immigrant was sent to the trachoma ward in one of the immigrant hospitals, and was admitted to the U.S.A. after treatment. The more likely outcome of the committee hearing was that the immigrant would be sent back to his port of origin.<sup>30, 49</sup> The U.S. government did not change the rules for special immigrants and detained two sons of a Chinese government official in 1923 at the Angel Island immigration station in California. Both men had trachoma and the U.S. government ignored their complaints that they deserved better treatment.<sup>19</sup>

**Trachoma was viewed as a serious threat by many governments and this threat guided decisions regarding immigration legislation.**

While steerage passengers encountered a rigorous inspection ashore, first and second-class travelers were inspected onboard the ship. Only those showing symptoms were sent with the steerage passengers to be inspected more thoroughly. An ill first or second class passenger had a much better chance of slipping through the inspections than an ill steerage class passenger. This became known as the second-class scam, and any immigrant who was able to purchase the more expensive ticket did so.<sup>40</sup> The medical exam was an involved process that created apprehension for the immigrants.

1897

Trachoma classified as a loathsome and contagious disease by the US

1902

Canada passes laws excluding immigrants with trachoma

1903

Sir Ernest Cassel donates £40,000 for combating eye disease in Egypt

1905

All immigrants entering the US must be examined for trachoma

### Trachoma and Immigration

The sanitation of the inspections at Ellis Island remains a debated topic. When Theodore Roosevelt visited in 1906 he commented that “the Line,” as the inspections were referred to, was unsanitary and appeared to be a way to spread disease to healthy people. The inspectors were supposed to wipe the buttonhooks with Lysol soaked towels and wash their hands between each inspection, yet, with limited time and facilities for sanitation, this did not always happen.<sup>41</sup> Fortunately, however, no outbreaks were reported among immigration inspectors or among newly arriving immigrants.<sup>31, 74</sup>




Buttonhooks were commonly used to evert immigrants' eyelids.

While the USPHS spent 80% of its resources on medical inspections at the borders, on average less than 1% of incoming immigrants suffered from trachoma each year.<sup>41</sup> However, trachoma did represent the leading medical cause for rejection of immigrants at the U.S. border.<sup>16, 19</sup> Although preventing immigrants from entering the country with trachoma certainly helped avoid new cases, immigrants were not the only people suffering from trachoma in the United States. The rates of trachoma were much higher than 1% in some areas of the United States. A survey in Kentucky in 1912 found more than 12% of those surveyed suffered from trachoma.<sup>3</sup> Trachoma rates were high in Virginia, Kentucky, Tennessee, Missouri, Arkansas, Alabama, and Oklahoma, which became known as the Trachoma Belt. Additionally, the prevalence of trachoma reached 90% on some Native American reservations, while surveys of the students in New York City public schools found that about 10% of students suffered severe or mild trachoma.<sup>3, 31, 41</sup>

### Immigrants were not the only people suffering from trachoma in the United States.

Despite the attention given to identifying cases of trachoma, little was known about effective treatments. Although no effective remedy was available, many different treatments were being administered. Those immigrants who were placed in hospitals were given a variety of therapies including removal of infected tissue, rupturing of the granules, rinsing the eye with different washes, and rubbing the eyelid with a “bluestone” which was made of copper sulfate crystals.<sup>40</sup> A large market for medications flourished, in spite of the fact that they were ineffective. Some of these treatments were aimed at curing the disease, while others were intended to cover the symptoms of trachoma in order to trick the immigration inspectors. San Francisco newspapers documented the topical use of adrenaline by Chinese and Japanese immigrants to mask the signs of trachoma and fool immigration officers.<sup>19</sup> Some immigrants also rubbed their eyelids with sugar cubes for six weeks before embarkation to cure themselves of trachoma symptoms.<sup>16</sup>



1906

Teddy Roosevelt visits Ellis Island and raises concern over inspection sanitation

Halberstaedter and Prowazek discover trachoma inclusion bodies

1907

First trachoma hospital opens in the US

President Wilson funds trachoma research

1913

MacCallan creates 4 stage classification of trachoma


1914

Portugal prohibits entrance of immigrants with trachoma

1923

Chinese diplomat's sons are detained for trachoma in US

Approximate geographical distribution of trachoma based on estimates by various authors survey data, information from eye clinics or notification of cases.



### Trachoma and Immigration

With immigrants and travelers moving around the world, health and disease control became something to address on a global level. Between World War I and World War II La Ligue Contre Le Trachome was organized and worked with the International Organization Against Trachoma, which was founded in 1929. Additionally, the World Health Organization was formed in 1948 and immediately identified trachoma as an important disease to combat. In 1949 the WHO collected all the available statistics on trachoma rates worldwide and put together a report. A summary of this information is found above.<sup>72</sup>

### The Discovery of *Chlamydia trachomatis* and Elimination of Trachoma

The twentieth century brought many scientific advances that aided efforts to eliminate trachoma. In 1907, Halberstaedter and Prowazek found cytoplasmic inclusion bodies, small structures in the cytoplasm of cells, in samples from apes and humans suffering from trachoma. As research continued, scientists discovered that these inclusion bodies were found in non-gonorrheal ophthalmia neonatorum, male urethritis, and female genital epithelium.

This began linking trachoma, newborn conjunctivitis, and non-gonorrheal sexually transmitted infections. Studies continued to determine the nature of the Halberstaedter-Prowazek inclusion bodies and around 1930 these inclusion bodies were accepted as the cause of trachoma.<sup>7, 16, 37, 60, 65</sup>

In 1954 T'ang and his colleagues in China cultured *Chlamydia trachomatis* in the yolk sac of chick eggs. In addition to T'ang, many other scientists were attempting to cultivate the agent of trachoma and were producing a variety of results. This prompted the WHO to create criteria for acceptance of an attempt as successful. T'ang's work met the guidelines and in 1957 his accomplishment was recognized. Throughout the first half of the 20th century, the agent of trachoma was thought of as a virus because it was so small and could only be grown within living cells. By the 1970's, however, *Chlamydia trachomatis* was acknowledged as a bacterium because it possesses both DNA and RNA and is susceptible to antibiotics.<sup>7, 16, 37, 60, 65</sup>

**Trachoma** is mentioned in Mario Puzo's *The Godfather*, in Bill Bryson's *In a Sunburned Country*, and in Mark Twain's *Innocents Abroad*

Doctors successfully treat trachoma with sulfonamides

1937

1940

First use of penicillin to treat trachoma

1943

First use of streptomycin to treat trachoma

Francis I Proctor Eye Research Labs open at UCSF

1947

## The Discovery of *Chlamydia Trachomatis* and Elimination of Trachoma

Along with identification of the agent came several new therapies for treating trachoma. The discovery of antimicrobials in the mid-20th century aided the progression of trachoma treatment even more than the identification of the agent.<sup>16</sup> In 1937, scientists discovered that sulfanilamide was effective for treating trachoma.<sup>3</sup> However, many people were allergic to sulfa antibiotics and experienced a severe skin reaction, so efforts continued to find an alternative treatment. In the early 1950's, both topical and oral tetracyclines were investigated. Oral or systemic tetracyclines were found effective, but slowed bone growth and damaged the teeth of young children, the primary group to receive therapy.<sup>17</sup> Topical tetracyclines were chosen as the most effective therapy with the fewest side effects. Topical tetracyclines remained the treatment of choice until the 1990's when oral azithromycin therapy, which requires only one annual dose, was developed and became the preferred treatment for trachoma. With these new therapies available, trachoma control became a more manageable task.

Several strains of *Chlamydia trachomatis* exist, some of which cause trachoma, while other genetically different strains cause a sexually transmitted infection. The research on *C. trachomatis* was initially useful for combating both diseases. In the 1970's and 1980's, however, the focus of research efforts shifted to the sexually transmitted infections. In 1973, work to eliminate trachoma by the WHO became part of the effort to prevent blindness, which gave trachoma importance in a larger context. Additionally, in the 1980's, the Edna McConnell Clark Foundation's (EMCF) continued support of research specifically targeting blinding trachoma was crucial to trachoma elimination efforts. Work funded by EMCF included the development by the WHO of a simplified scheme for diagnosing trachoma in the field, research investigating risk factors associated with trachoma, and the establishment of the value and feasibility of surgery.<sup>8</sup>

EMCF also worked to advance the comprehensive SAFE strategy, which is the basis for elimination efforts today. The SAFE strategy combines treatment (surgery and antibiotics) and prevention (face washing and environmental improvement). While the separate components of the SAFE strategy had been used throughout history, the packaging of the components into a four-part strategy offered a comprehensive approach to trachoma elimination that was adaptable to many different situations and could be implemented at the community level.



Face washing is an important component of the SAFE strategy. Just 1 liter of water can clean up to 30 children's faces.

In 1996, the WHO, with support from EMCF, convened a global scientific meeting on trachoma. At the time of the meeting scientists had established that azithromycin was going to change trachoma control efforts. Participants at the meeting "identified 46 countries as having known areas of blinding trachoma" and created a list of 16 priority countries for trachoma control.<sup>73</sup> The following year, the WHO established the GET (Global Elimination of Trachoma) 2020 program in cooperation with national health programs, non-governmental organizations, research institutions and other partners, many of whom attended the scientific meeting in 1996. The GET 2020 alliance endorsed the SAFE strategy and has been working in endemic countries around the world to eliminate blinding trachoma with the goal of elimination by the year 2020. The GET 2020 alliance falls within the VISION 2020 program, which began work in 2000. These steps taken by the WHO aided the visibility and cooperation of the elimination efforts.

First WHO trachoma expert committee meeting

1949

First WHO assembly recognizes need for trachoma control

1952

1954


Chinese culture trachoma agent

1957

Isolation of agent accepted by WHO

Establishment of the International Association for the Prevention of Blindness

1975



### The Discovery of *Chlamydia Trachomatis* and Elimination of Trachoma

The recognition of the effectiveness of a single, oral dose of azithromycin for the antibiotic component of the SAFE strategy was crucial to trachoma elimination progress. Pfizer's antibiotic Zithromax® (azithromycin), which is produced and used to treat many bacterial infections, was found effective for treating trachoma in clinical trials. Through the efforts of EMCF, Pfizer committed to join the fight against trachoma by donating both Zithromax® and financial support to the International Trachoma Initiative (ITI), an organization formed by EMCF and Pfizer in 1998 to support the control efforts. As the only organization dedicated solely to trachoma, ITI has played an important role in working towards the goal of global elimination. The work completed by EMCF also motivated many other organizations to undertake trachoma control projects.

Today, national ministries of health, non-governmental organizations, and other partners are working together to ensure that surgical services are available to patients with advanced disease, antibiotics are distributed, face washing is widely publicized, and communities are working to improve access to clean water and sanitation. The goal set by the WHO is elimination of blinding trachoma by 2020, although some countries, such as Morocco, may achieve this goal much sooner.<sup>27</sup>



Children in Tanzania are educated about the four components of the SAFE strategy

1978

Establishment of WHO Program for the Prevention of Blindness

EMCF publishes trachoma research plan

1985

1987

WHO adopts simplified grading scheme

1991


First discussions with Pfizer and EMCF



**For**  
the individuals suffering from advanced trachoma

the disease has always been devastating, but with the continued success of ITI and other organizations, it is hoped that the long and complex history of blinding trachoma will soon come to a close. Trachoma disappeared from the United States and Europe with improved sanitation and limited public health efforts. People around the world, with help from the international community, can be free from blinding trachoma without waiting for the disease to disappear on its own. With elimination of blinding trachoma, millions of people will be spared the painful and disabling path towards blindness. Additionally, helping individuals retain their sight for the duration of adulthood will allow them to be more productive, and will help break the cycle of poverty. With hard work and continued commitment, the world can be free of the burden of blinding trachoma.

**People** around the world, with help from the international community, can be free from blinding **trachoma** without waiting for the disease to disappear on its own.



## References

- Al Arab, Gamal Ezz, Nagah Tawfik, Reda El Gendy, Wagida Anwar, and Paul Courtright. "The burden of trachoma in the rural Nile Delta of Egypt: a survey of Menofiya governorate." *British Journal of Ophthalmology* 85 (2001): 1406-1410.
- Albert, Daniel M. and Diane D. Edwards ed. *The History of Ophthalmology*. Cambridge, MA: Blackwell Science, 1996.
- Allen, Shannen K. and Richard D. Semba. "The Trachoma 'Menace' in the United States, 1897-1960." *Survey of Ophthalmology* 47.5 (2002): 500-508.
- Al-Rifai, Ikassim M. J. "Trachoma through history." *International Ophthalmology* 12.1 (1988): 9-14.
- Aristophanes. *Plutus*, 22 July 2004 <[http://www.greece.com/library/aristophanes/plutus\\_01.html](http://www.greece.com/library/aristophanes/plutus_01.html)>.
- Aronson, Stanley M. "Med-school dropout in Renaissance Italy." *Providence Journal-Bulletin* 22 December 2003: A-15.
- Arrington, George E. Jr. A *History of Ophthalmology*. New York: MD Publications, Inc., 1959.
- Asbury, Carolyn, Barnett L. Cline, and Victoria M. Gammino. "The Edna McConnell Clark Foundation's Tropical Disease Research Program: A 25-Year Retrospective Review 1974-1999." 15 Aug. 2000. <<http://www.emcf.org/about/pastprograms/iti.htm>> 21 July 2004.
- Baasanhu, J. and G.J. Johnson, G. Burendei, and D.C. Minassian. "Prevalence and causes of blindness and visual impairment in Mongolia: a survey of populations aged 40 years and older." *Bulletin of the World Health Organization* 72.5 (1994): 771-776.
- Bietti, G.B. "Natural History and Diagnosis of Trachoma." *Israel Journal of Medical Science* 8.8 (1972): 1101-1106.
- Boldt, J. *Trachoma*. London: Hodder and Stoughton, 1904.
- Cong. *An act in amendment to the various acts relative to immigration and the importation of aliens under contract or agreement to perform labor*. US 51st Cong., 2nd sess. Chap. 551. 3 Mar. 1891. 26 July 2004 <[http://web.lexis-nexis.com/congcomp/attachment/a.pdf?\\_m=ac0cf1680204fd0449481c4074cd81cb&wchp=dGLbVtb-zSkSA&\\_md5=c5a7eb43fbd989ae789781c01a753920&ie=a.pdf&gt;](http://web.lexis-nexis.com/congcomp/attachment/a.pdf?_m=ac0cf1680204fd0449481c4074cd81cb&wchp=dGLbVtb-zSkSA&_md5=c5a7eb43fbd989ae789781c01a753920&ie=a.pdf&gt;)>
- Conway, Lorie. "The Other Ellis Island: The Untold Story of the Ellis Island Hospital." A Documentary Film Proposal. Ellis Island Library Archives.
- Cook, Joseph. Timeline of founding of ITI. Personal documents.
- Crowe, Sam. "The Maasi move to eliminate trachoma." *British Medical Journal* 314.7097 (1997): 1781.
- Dawson, Chandler R. Telephone interview. 24 June 2004. and 10 August 2004.
- Dawson, Chandler R., and Julius Schacter. *Human Chlamydial Infections*. Littleton, Mass.: PSG Pub. Co., 1978.
- Dawson, C.R., T. Daghfous, I. Hoshiwara, K. Ramdhane, M. Kamoun, C. Yoneda and J. Schacter. "Trachoma therapy with topical tetracycline and oral erythromycin: a comparative trial." *Bulletin of the World Health Organization* 60.3 (1982) 347-355.
- Fairchild, Amy L. *Science at the Borders: Immigrant Medical Inspection and the Shaping of the Modern Industrial Labor Force*. Baltimore: Johns Hopkins University Press, 2003.
- Fairchild, Amy. "Policies of Inclusion." *American Journal of Public Health* 94.4 (2004): 528-539.
- Feibel, Robert M. "John Vetch and the Egyptian Ophthalmia." *Survey of Ophthalmology* 28.2 (1983): 128-134.
- Frank, Al. "Midwives to America." *The Star-Ledger* 29 September 1989.
- Frick, Kevin. Telephone interview. 5 July 2004.
- Frick, Kevin D., Eva Basilion, Christy Hanson, and M. A. Colchero. "Estimating the burden and economic impact of trachomatous loss." *Ophthalmic Epidemiology* 10.2 (2003): 121-132.
- Ho, Viet H. and I. R. Schwab. "Social Economic Development in the Prevention of Global Blindness." *British Journal of Ophthalmology* 85 (2001): 653-657.
- "International League Against Trachoma." *The Lancet* 216.5581 (1930): 359.
- International Trachoma Initiative Dedicated to eliminating the world's leading cause of preventable blindness*. 2002. International Trachoma Initiative. 21 June 2004 <<http://www.trachoma.org>>.
- Jones, Barrie R. "The prevention of blindness from trachoma." *Transactions of the Ophthalmological Society of the United Kingdom: The Bowman Lecture, 1975* 95 (1975): 16-32.
- Karasch, Mary C. "Ophthalmia." *The Cambridge World History of Human Disease*. Cambridge University Press. (1993): 897-905.
- Kraut, Alan M. "Germs, Genes, and American Efficiency." *Social Science History* 12.4 (1988): 377-388.
- Kraut, Alan M. *Silent Travelers: Germs, Genes and the "Immigrant Menace"*. New York: Basic Books, 1999.
- Kuper, Hannah, A. W. Solomon, J. Buchan, M. Zondervan, A. Foster and D. Mabey. "A critical review of the SAFE strategy for the prevention of blinding trachoma." *The Lancet Infectious Disease* 3 (2003): 372-379.
- Kupka, K., B. Nizetic and J. Reinhardt. "Sampling studies on the epidemiology and control of trachoma in southern Morocco." *Bulletin of the World Health Organization* 39.4 (1968): 547-566.
- Kyaw, T. A., T. Nyunt, T. K. Sundaresan, and M. L. Tarizzo. "Control of trachoma and prevention of blindness in rural communities in Burma." *Bulletin of the World Health Organization* 56.6 (1978): 945-955.
- Larner, A. "Ophthalmological Observations Made During the Mid-19th-century European Encounter with Africa." *Archives of Ophthalmology* 122.2 (2004): 267-272.
- Lavery, Francis S. "Trachoma in Ireland." *The Lancet* 216.5595 (1930): 1163-1166.
- Mabey, David and R. Bailey. "Eradication of trachoma worldwide." *British Journal of Ophthalmology* 83.11 (1999): 1261-1263.
- MacCallan, A. F. "The Relationship Between Conjunctivitis and Trachoma." *General Assembly of the International Association for the Prevention of Blindness* Paris, 1936.
- MacCallan, A. F. *Trachoma*. London: Butterworth: 1936.
- Markel, Howard. "'The Eyes Have It': Trachoma, the Perception of Disease, the United States Public Health Service, and the American Jewish Immigration Experience, 1897-1924." *Bulletin of the History of Medicine* 74 (2000): 525-560.
- Markel, Howard. *When Germs Travel: Six Major Epidemics That Have Invaded America Since 1900 and the Fears They Have Unleashed*. New York: Pantheon, 2004.
- Mecaskey, Jeffrey W., Charles A. Knirsch, Jacob A. Kumaresan, and Joseph A. Cook. "The possibility of eliminating blinding trachoma." *The Lancet Infectious Disease* 3.11 (2003): 728-734.
- Meredith, Suzanne, J. Headley, G. Peach and Dayalan Devanesen. "Trachoma in the Northern Territory of Australia, 1940-1986." *The Medical Journal of Australia* 151 (1989): 190-196.
- Millar, Marcia Inhorn and Sandra D. Lane. "Ethno-Ophthalmology in the Egyptian Delta: An Historical Systems Approach to Ethnomedicine in the Middle East." *Social Science & Medicine* 26.6 (1988): 651-657.
- Mullan, Fitzhugh. *Plagues and Politics the Story of the United States Public Health Service*. New York: Basic Books Inc., 1989.
- Munoz, B. and S. West. "Trachoma: the forgotten cause of blindness." *Epidemiologic Reviews* 19.2 (1997): 205-217.
- Pizzarello, Louis et. al. "VISION 2020: The Right to Sight: A Global Initiative to Eliminate Avoidable Blindness." *Archives of Ophthalmology* 122.4 (2004): 615-620.
- Prost, A. and A. D. Negrel. "Water, trachoma and conjunctivitis." *Bulletin of the World Health Organization* 67.1 (1989): 9-18.
- Reed, Alfred C. "The Medical Side of Immigration." *Popular Science Monthly* April (1912): 383-392.
- Resnikoff, Serge and Ramachandra Pararajasegaram. "Blindness Prevention Programs: Past, Present, and Future." *Bulletin of the World Health Organization* 79.3 (2001): 222-226.
- Savage-Smith, E. "Ibn al-Nafis's Perfected book on ophthalmology and his treatment of trachoma and its sequelae." *Journal for the History of Arabic Science* 4 (1980): 147-204.

## References

52. Schemann, J. F., D. Sacko, A. Banou, S. Bamani, B. Bore, S. Coulibaly, and M. A. el Mouchtahi. "Cartography of trachoma in Mali: results of a national survey." Bulletin of the World Health Organization 76.6 (1998): 599-606.
53. Strategic Plan for Trachoma Research April 1990. Edna McConnell Clark Foundation.
54. Tabbara, K. F. "Blindness in the Eastern Mediterranean countries." British Journal of Ophthalmology 85.7 (2001): 771-775.
55. Taborisky, J. "Historic and Ethnologic Factors in the Distribution of Trachoma." American Journal of Ophthalmology 35 (1952): 1305-1311.
56. Tarrizzo, Mario. "The World Health Organization and the Prevention of Blindness." Transactions American Academy of Ophthalmology and Otolaryngology 79 (1975): 453-456.
57. Taylor, Hugh R. "A trachoma perspective." Ophthalmic Epidemiology 8.2 (2001): 69-72.
58. The Bible Online. Gen. 29:17. 22 Jul. 2004 <<http://www.bible.com>>.
59. "The Problem of Trachoma." The Lancet 216.5595 (1930): 1140.
60. Thygeson, P. "Trachoma virus: historical background and review of isolates." Annals of the New York Academy of Sciences 98 (1962): 6-13.
61. Thylefors, B. "Global Trachoma Control- Past, Present and Future Approaches." Revue internationale du trachoma et de pathologie oculaire tropicale et subtropicale et de sante publique 72 (1995): 13-80.
62. Thylefors, B., A. D. Negrel, R. Pararajasegaram and K. Y. Dadzie. "Global Data on Blindness." Bulletin of the World Health Organization 73.1 (1995): 115-121.
63. Tower, Paul. "The History of Trachoma." Archives of Ophthalmology 69 (1963): 157-164.
64. Treharne, John D. "The Community Epidemiology of Trachoma." Review of Infectious Diseases 7.6 (1985): 760-764.
65. Wang, Y. "Etiology of Trachoma: A great success in isolating and cultivating Chlamydia Trachomatis." Chinese Medical Journal 112.10 (1999): 938-941.
66. Watts, Susan. "The History of Trachoma in Egypt." Report for the International Trachoma Initiative (Unpublished) March 2003.
67. Webb, Stephen. "Prehistoric Eye Disease (Trachoma?) in Australian Aborigines." American Journal of Physical Anthropology 81.1 (1990): 91-100.
68. Wheeler, Maynard C. The American Ophthalmological Society The First Hundred Years. Toronto: University of Toronto Press, 1964.
69. Wilson, Rowland P. "The Various Forms of Conjunctivitis Among Children in Egypt; Their Seasonal Incidence and their Influence on the Development of Trachoma." General Assembly of the International Association for the Prevention of Blindness Paris, 1936.
70. World Health Organization Expert Committee on Trachoma. "Expert Committee on Trachoma: first report [of a meeting held in Geneva from 3 to 8 March 1952]." Geneva: World Health Organization, 1952.
71. World Health Organization. "Blindness as a public health problem in China." Fact Sheet No. 230. 15 July 2004. Sept. 1999 <<http://www.who.int/mediacentre/factsheets/fs230/en>>
72. World Health Organization. "World Distribution and prevalence of trachoma in recent years." Epidemiological and Vital Statistics Report Nov-Dec Geneva 1949.
73. World Health Organization. Programme for the Prevention of Blindness and Deafness. Future approaches to trachoma control: report of a global scientific meeting. Geneva, 17-20 June 1996. Geneva: World Health Organization, 1997.
74. Yew, Elizabeth. "Medicine, Bureaucracy and Immigration: Medical Inspection of immigrants at Ellis Island 1891-1924." (Unpublished Manuscript) Ellis Island Library Archives.
75. Zouboulis, C. "Benediktos Adamantiades and his forgotten contributions to medicine." European Journal of Dermatology 12.5 (2002): 471-474.

[www.trachoma.org](http://www.trachoma.org)



international  
**trachoma**  
initiative

441 Lexington Avenue  
Suite 1101  
New York, NY 10017  
T 212.490.6460  
F 212.490.6461

