An introduction to

THE NATURAL HISTORY OF PARADISE, by Karla Tanner, rev. 1997

THE SETTING

A. Identification

B. Climatic influences

C. Challenges to survival

1. Length of season
2. Volcanic soil
3. Wind
4. Ultraviolet radiation

I. **IDENTIFICATION and SURVIVAL STRATAGIES**

SURVIVAL STRATEGIES

A. Trees
1. Subalpine fir
2. Mountain Hemlock.

B. Shrubs
1. Sitka mountain-ash
2. Cascades blueberry
3. Rosy spiraea
4. Pink heather
5. White heather

C. Flowers
1. Avalanche lily
2. Glacier lily
3. Western pasqueflower
4. Snow cinqueflower
5. Subalpine buttercup
6. Sitka valerian
7. Meadow parsley
   a. Hoverfly
8. Broadleaf lupine
9. Dwarf lupine
   a. Bumblebee
10. Magenta paintbrush
11. Marshmarigold
12. Jeffrey shootingstar
13. Lewis monkeyflower
14. Beargrass
15. Tolmie saxifrage
16. Spreading phlox
17. Partridgefoot
18. Mountain bistort
19. Corn lily
20. Mountain daisy
21. Fireweed
22. Mountain bog gentian

D. Sedges
1. Black alpine sedge
2. Showy sedge

E. Lichen
1. Jewel lichen
2. Goat's beard lichen

F. Birds
1. Rosy finch
2. Rufus hummingbird
3. White-tailed ptarmigan
4. Blue grouse
5. Varied thrush
6. Gray Jay
7. Clark's nutcracker
8. Dark-eyed junco
9. Common Raven

G. Mammals
1. Yellow Pine chipmunk
2. Golden-mantled ground squirrel
3. Hoary marmot
4. Pika
5. Snowshoe hare
6. Black bear
7. Black-tailed deer
8. Red fox

III. **REFERENCES**

THE SETTING

It is often said that traveling form the Nisqually Entrance to Paradise is the same as traveling from Puget Sound to the Arctic Circle in a straight line. In general a difference of 1000 feet in elevation is equivalent to a difference of 300 miles in latitude or three degrees in temperature.

If temperature and moisture were uniform at a given altitude, the boundaries of where plants and animals could survive would be as distinct as contour lines on a topographic map. However, there is no distinct line of demarcation and there is often considerable variation in the altitudinal distribution of plants. On northern exposures, for example, the lack of sunshine results in lower temperatures and higher soil moisture. Plants which normally occur at 5000 feet on a sunny southern exposure may be found only at a lower elevation on northern slopes, and vise-versa.

Such variation is even more marked in the distribution of mammals and birds. Many species are characteristic of more than one are in relation to the season of the year, the scarcity or abundance of food, or other factors.

Paradise is considered to be a subalpine environment. THE CLIMATE is characterized by short, cool growing seasons and long snowy winters. Snow flurries may come at anytime during the year but generally snow starts staying on the ground starting in mid-October. The greatest snowpack is generally 15 feet to 20 feet deep and occurs during February or March. The weather station located along Nisqually Vista Trail measures an average of 630 inches of snow fall during the snow year of July 1- June30, with snow often staying on the ground there until the first week of July. Elsewhere around Paradise snow might take much longer to melt than early July, especially on north-facing slopes or in valleys where it accumulates to a greater depth. During the winter of 1971-72, 1122 inches (93.5 feet) of snow fell at Paradise – almost twice as much as the average. Two winters later, over 1000 inches of snow again blanketed the meadow. The snowpack was correspondently deeper than usual during those years and cooler areas of the meadow never completely melted out during the summers which followed.

Record amounts of snow fall at Paradise due to the height of Mount Rainier and due to the mountain’s close proximity to the ocean; as air masses blow off the Pacific, they are heavily laden with moisture. Mount Rainier is the first major barrier encountered by these eastward-flowing storms. As the air masses rise in order to move over the top of the mountain and continue their journey across the continent, the air cools. Since cooler air can hold less moisture, the excess moisture condenses in the form of rain or snow.

Because the storms are so heavily laden with moisture when they reach Mount Rainier, the snow which falls on the slopes of this mountain is typically much wetter than snow which falls further inland. Wetter snow takes longer to melt. Consequently, snow lingers at Paradise long after a

comparable depth of snow has melted in the Rocky Mountains which are much further away from the Pacific Ocean.

The LONG WINTER, coupled WITH A SHORT SNOW-FREE SUMMER growing season is the primary challenge facing plants trying to grow at Paradise.

The second major challenge is the soil. VOLCANIC SOIL is one of the most difficult soils for plant life. It contains little nourishment. The soil is as hard as ground glass to the touch and as gritty. It retains its separateness as individual grains even when wet. Volcanic soil drains so quickly that humus has almost no chance to form in the cool alpine climate. Vegetation has a hard time absorbing moisture before it is lost. Volcanic soil is also extremely abrasive as it contains large amounts of bare rock and rubble. As it dries, it floats away with a breath of wind.

WIND is the third most important challenge. It is a constant presence. Not only does wind lower temperatures by the severity of the chill factor – a 30 mph wind can lower the temperature 20 degrees – but wind velocity also erodes soil and increases evaporation. Wind dictates the shape of plants, scatters seeds and pollen, and molds the snow surface into fanciful sculptures. When wind is strong, it carries sand and snow and has a considerable abrasive effect, sandblasting bark and twigs on the windward side of trees. If water is not available to trees of other plants at these times, either because the soil is frozen or because moisture is lacking, strong ground winds and cause plants to wilt as if under severe drought. Wind desiccation is most lethal in winter when it prunes all elevated growth.

A further consideration is local up slope winds which are caused by ground heating in the lowland . The air at lower elevations absorbs ground warmth from the sun-heated soil; the air expands, rises and cools. At sunset, when the soil on high mountain ridges cools rapidly, the thin air loses its heat quickly and becomes heavier, flowing down the slope at night. Cold air drainage at night makes ridges warmer than the bottom of subalpine valleys. Plants and animals are unable to survive in cold valleys may find more favorable living conditions on warmer ridges. One need only stand near Myrtle Falls and look toward the Mountain to notice the lack of shrubs and trees in the Edith Creek Basin by comparison to the higher ridges all around.

And lastly, ULTRAVIOLET RADIATION and intense light are the fourth major challenges facing vegetation growing on the upper slopes of Mount Rainier. At 14,000 feet, for example, there is twice as much ultraviolet radiation and 25% more light than at sea level.

A short summer growing season, porous volcanic soil, wind, and intense light – these are the major factors to which vegetation of the subalpine and alpine zones on Mount Rainier must adapt in order to survive. *How* they have met the challenge of living high on the slopes of Mount Rainier is fascinating.

IDENTIFICATION AND SURVIVAL STRAGEGIES

TREES

The most abundant tree at Paradise, SUBALPINE FIR, is more characteristic of warmer, drier slopes. It can be identified by its rigid, cathedral-spire growth shape or pointy top and stubby upper branches. Although stiff, the branches do not break under the weight of snow since they offer only a small surface on which snow can accumulate. During winter, the longer lower branches are protected from branch-breaking snowfall, cold and desiccating wind by the snowpack.



Alpine fir, NPS photo

Although subalpine firs produce viable seeds the growing season at Paradise is often too short, cool and wet for the seeds to survive. (Conversely, at Sunrise, the summer is usually too hot and dry for the seeds to survive). For the most part these trees spread by *layering*, i.e., new roots and stems sprout from branches lying on soil beneath the parent tree. Of the tree species characteristic of Paradise, subalpine fir is the best at layering.

Layering benefits these trees in several ways: young trees share an established root system with the parent tree; as a group of trees becomes established they begin to exert an influence on their micro environment, i.e. lengthening the growing season because of their black body effect which results in earlier snowmelt; a group of trees provides some protection from wind velocity and desiccation for its individual members; and as wind is slowed through the clump it releases wind-blown soil particles.

In order for tree species to produce cellulose (wood), the temperature must average 50 degrees F for approximately 30 days. So, although treeline on this mountain is generally considered to be about 7000 feet elevation, trees will grow higher on warmer ridges than in valleys where colder air collects. Although temperatures might be more favorable on ridges, living in such exposed locations can be particularly difficult. Harsh drying winds, carrying sharp ice crystals, kills and cut away shoots from the windward side of trees while growth on the leeward side of these trees goes on normally, producing the one-sided appearance known as *flagging.* Height will be stunted when the trees grow higher than the insulating snow pack and the leader is trimmed off by the wind action leaving a skirt of growth below a gnarled top. These stunted conifers are sometimes referred to as *krummholz.* Subalpine fir tends to be more common in krummholz formations than other species.

Like all conifers, the needles of the subalpine fir are waxy thus reducing water loss through transpiration. The needles are also edible. Although snowshoe hares prefer tenderer flower stems and sedges when they are available during summer, the hares' diet during winter consists of readily-available needles from conifer trees.

Also successful at layering, MOUNTAIN HEMLOCK is the second most abundant tree at Paradise. It is characteristic of wetter locations and can be easily identified by its drooping leader and long, wispy branches. Even though the branches of the hemlock are long they are also quite flexible. Instead of breaking when snow loads accumulate, the branches flex, lose their snow and bounce back up. Hemlock trees produce short, waxy needles which radiate from the branches in a star-like pattern. As on all conifers, evergreen needles of the mountain hemlock are advantageous in an area where the growing season is short, as it is more energy efficient to use one set of leaves for more than a single season and to replace those leaves individually rather than all at once.

Mountain hemlock,
©2012 Gary A. Monroe

YELLOW CEDAR, PACIFIC SILVER FIR and a very occasional WHITE BARK PINE also grow at Paradise.

Yellow cedar
Connecticut Agricultural Experiment Station Archive

Pacific silver fir
Joseph Dougherty, M.D./ecology.org

White bark pine
Robert Potts © California Academy of Sciences

Climatic conditions on the west side of the park have generally been favorable for tree establishment since the close of the Little Ice Age (late 1800s). Warm, dry summers favor the survival of tree seedlings. In the 1960s park managers noticed large numbers of saplings in the meadow. Dr. Jerry Franklin conducted research into tree establishment patterns. He determined that tree establishment had increased between 1915 and 1940 during warm, dry periods. Subsequent research at Mount Rainier has demonstrated that tree establishment has continued until the present.

SHRUBS

One popular story about the SITKA MOUNTAIN-ASH is that the average depth of the winter snowpack can be told by the height of this shrub. Certainly the snow depth is now lower than its topmost branches. Snow insulates the shrub from winter cold and protects it from drying winter winds. Exposed branches do not

Sitka mountain ash, Barry Breckling

survive the winter. Until covered by snow, mountain-ash berries remain on the bush where the ferment as autumn progresses, causing some observers to report birds “flying under the influence” after feasting on fermented berries. Although they are rather mealy textured the bright orange clusters of berries are enjoyed well enough by birds such as varied thrush and by small mammals such as chipmunks and ground squirrels.

Huckleberry or blueberry? In the Pacific Northwest the two terms are used interchangeably. Of the 12 species of blueberries which inhabit this region, the CASCADES BLUEBERRY grows at the highest elevation. During summer the tiny bell-shaped flowers of these low-lying bushes are overshadowed by the more colorful blooms of other subalpine plants. But by October blueberry bushes provide 90% of the red-bronze color characteristic of autumn at Paradise. When the berries ripen at Paradise, black bears are more frequently seen in the meadows than earlier in the summer when foraging doesn’t yield as tasty of results.

Blueberry, Jean Pawek

A myriad of tiny blossoms cover the flowerheads of the shrubby plant named ROSY SPIRAEA. The stamens which stick up from the flowerheads give them a fuzzy appearance and tickle your nose when you try to smell them. The dense foliage provides shelter for many small mammals and birds such as rabbits and grouse.

Rosy spirea, JW Stockert; 1973

Pink heather, NPS Photo

Above Paradise where heather grows on the fellfields of Muir Snowfield, botanists have found these shrubby, mat-forming plant communities dating to a remarkable 7000 years. PINK HEATHER can also be found throughout the subalpine parkland where it generally favors stony soils. The leaves of this heather are waxy to help prevent wind desiccation. Like most flowers of subalpine and alpine environments, this heather is a perennial. It doesn’t have to expend valuable energy producing stems and leaves every year but simply adds to what is already established.

WHITE HEATHER favors windy areas above timberline where it forms mats and nestles among the rocks for warmth and wind protection. Unlike the needle-like leaves of other mountain heathers, the evergreen leaves of white heather for tiers of overlapping scales resembling clubmoss. White heather is credited with being naturalist John Muir’s favorite flower.

White heather, NPS Photo

FLOWERS

AVALANCHE LILY and GLACIER LILY base their success on timing rather than brute size. They begin growing their bulb tips in September, just when their neighbors are dying back. While snow covers these plants for the next eight months or so, holding the soil temperature close to freezing, their shoots inch up to the soil surface. Very few plant species are active at such low temperatures. Without the snow blanket temperatures would be even colder and growth would be impossible. As soon as the snow melts away from the shoots in spring, avalanche lily and glacier lily burst to their full heights, expending in a few days their disproportionately large reserve of starches. The stems of these little plants can push through the last inch or two of snow by combusting starch to melt a hole in the snow through which they grow. A remarkable adaptation in these and many of the early-blooming high-country plants is a thin-fleshed hollow stem used as an internal greenhouse. When the stored carbohydrates are burned off during the quick burst of growth, some of the heat produced is retained in the stem, making the internal air temperature warm enough for photosynthesis even when the outside air is not. Waste carbon dioxide from respiration also stay inside, available for synthesis into new carbohydrates. Although glacier lilies favor dry, south-facing slopes and generally only bloom early in the summer, avalanche lilies bloom at Paradise from late may through early August. As summer progresses, look for avalanche lilies along the edges of retreating snowbanks in meadow depressions where snow accumulation is greater, or on shady north-facing slopes, or in cooler alpine elevations. During wet weather, the flowers of both avalanche and glacier lilies droop to protect pollen inside the blossoms.

Avalanche lily, NPS Photo

Like avalanche and glacier lilies, WESTERN PASQUEFLOWER is one of the first flowers to bloom at Paradise. A “fur coat” of fuzzy hair helps protect more delicate areas of the plant. The hairs diffuse the strong alpine light, reducing both the intensity and directness of that reaching the surface of the leaves and stem. Although by itself the intensity of the light is not sufficient to upset photosynthesis, it may cause cell damage if combined with high reflection from rock or snow surfaces. The same hairs which protect the plant from excessive light also measurably reduce water loss by keeping drying winds off the plant’s stems and leaves, by encouraging condensation of moisture from the air, and by maintaining warmth. Pasqueflower blossoms are heliotropic, meaning they track the sun’s course through the day in order to maximize their exposure to light and warmth. As solar heat receptors in a yet snow-covered meadow, the warm pasqueflower blossoms make an appealing landing zone for pollinators. When flowering in early summer, the plant is only about 5 inches tall. By the time it fruits at the end of summer it has grown to nearly 24 inches tall and looks much like a mop head. Since the seeds are wind

Western anemone, NPS Photo

distributed this additional height allow them to be blown further from the parent plant. Pasqueflower is also commonly known as Western anemone.

Cinquefoil, NPS Photo

Although often confused with buttercups, cinquefoils can be identified by remembering their common name originated from the words meaning *five* and *leaf*. Cinquefoil's five yellow petals alternate with their five green sepals which alternate with their five shorter green bracts. The underside of SNOW CINQUEFOIL leaf is covered with crinkled hairs, looking like a wool ski sock under the microscope. Another way to distinguish the two plants is to look for the peculiar waxy sheen of the SUBALPINE BUTTERCUP's yellow petals. This sheen reflects away some of the sun's radiant heat—a valuable asset for a plant known as a herald of spring. The glossy surface of the buttercup petals also attracts inexperienced insect visitors, such as flies, more effectively than a matte surface. Buttercups start to grow while still under the snow and bloom in the meltwater of receding snowbanks.

Valerian, NPS Photo

As subalpine meadows are released by spring snowmelt, deep reddish shoots of SITKA VALERIAN soon shoot up abundantly. The redness disappears as the foliage matures, lingering longest in the budding flowers but disappearing as they mature to white. Most redness in plants comes from anthocyanin, a complex carbohydrate pigment that may be red, blue, or anywhere in between. Anthocyanin is suspected of several functions in high-elevation plants. First, it filters out ultraviolet radiation, which can be at least as hard on plant tissue as on human skin. Ultraviolet is especially intense at high elevation, where there is less atmosphere to screen it, and around the solstice in June, when sunlight is at its peak. In June the high country is still snowbank-chilled, and plant tissues young and tender, so that's where and when anthocyanin is brought out. Second, while reflecting ultraviolet light radiation, it also seems to absorb and concentrate infrared, thus heating the plant. Third, anthocyanin is an interim form for carbohydrates on their way up from winter storage. In order to bloom and fruit early in their short growing season, high-country plants store carbohydrates in their roots, and then move them up fast after snowmelt, or even before. In de-reddening, valerian stuffs itself with preserves from the root cellar. Its sweet smell and sturdy flattish cluster of tiny white blossoms make the valerian a good flower for pollination by butterflies. Since butterflies do not hover, they use the flower head as a landing pad on which to rest and open their wings while they sip nectar.

Meadow parsley, NPS Photo

Although sometimes confused with Sitka valerian or Queen Anne's lace, MEADOW PARSLEY can be identified by its open, round-topped clusters of tiny white-to-pinkish flowers. Its minute flowers

cluster together in a flat umbrella-like head and are favorites of hover flies. As pollinators, HOVER FLIES come in second only to bees. Flies are especially crucial in high mountain meadows, where bumble bees are the only common bees. Flies mimic the black-and-yellow-banded color pattern of bees and wasps as protection against any predator that made the mistake of attacking a bee or wasp. Hover flies even mimic the sound of bees, employing similar wingbeat frequencies. Hover flies, however, have a slender, non-furry body. Flowers preferred by flies are usually regular, simple, and often open flat, like parsleys and buttercups. Nectar guides are often present and nectar is easily available. Unlike bee flowers, fly flowers usually produce little odor.

Broadleaf lupine, NPS Photo

Since lupines can often be seen growing in areas where other plants cannot, it was once believed they devoured all of the nutrients from the soil. The plant received a name meaning “wolf” as a result. However, we now know that lupine, like all members of the legume family, fertilize the soil as they grow by releasing nitrogen from nodules on their roots. As the soil is improved, other plants can move into the area to grow alongside the lupines. The shee apparent on the leaves of the BROADLEAF LUPINE reflects away some of the sun’s radiant heat, while the hairs keep drying winds off moist surfaces. Alpine or DWARF LUPINE is even more covered with downy hair which gives the plant a soft, pale grey-green appearance. The hairs protect the plant’s stomata, the pore through which a plant breathes and represents an environmental response for conserving moisture. When moisture condenses on the leaf, it is directed by the configuration of the leaf whorl toward the center stem whee a drop of water can frequently be found long after all moisture has long evaporated from neighboring flowers. Lupine flowers produce a sweet scent that can be especially strong on warm days. The perfumey scent serves to attract insects such as mosquitos and bumble bees as pollinators. Bumble bees prefer blue flowers, the color of most lupine flowers at Paradise.

Bumble bee, NPS Photo

Bees are the most valuable pollinators of the plant world. BUMBLE BEES, the most common bees, are in the honey bee family, and have a strong social order though their colonies are relatively small and short-lived. Only the queens eat well enough to survive the onset of winter. The rest of the colony dies in the fall, after the drones and queens have mated. By feeding only a small number of queens per colony in preparation for hibernation, the colony conserves nectar and pollen resources which are scarce in cool climates. Other key adaptations include the use of ready-made insulated nests, typically a mouse or vole burrow; the relatively large, furry bodies and the skill of raising a near-constant body temperature for flight in a wide range of weather conditions, e.g., arctic bumble bees have been seen in flight in a snowstorm at 6 degrees below freezing!

What appear to be bright rose-purple flowers on MAGENTA PAINTBRUSH are really modified leaves called *bracts*. The colorful bracts are designed to attract pollinators such as the Rufus hummingbird to the nectar produced by tiny elongated flowers situated on the top of the bracts along the stem. Paintbrush is one of the several members of the figwort family known to parasitize the roots of other plants. This increases the amount of moisture available to the paintbrush since it virtually steals moisture away from its neighbors. Paintbrush benefits from an abundance of hairy fuzz in the same way as do its neighboring plants such as lupine.

Paintbrush, NPS/ Chris Roundtree

Not all the plants of the Hudsonian Zone have ways to conserve moisture. Those plants that do not, adapt to life in the subalpine meadows by growing only in areas with a readily available, constant source of moisture. MARSHMARIGOLDs can be found growing in the meltwater of snowbanks as early as ay on some southfacing slopes at Paradise. As the summer progresses, they are often seen blooming alongside JEFFERY SHOOTINGSTAR in meadow marshes and along the banks of streams, ponds, and wet seepages. And by late summer LEWIS MONKEYFLOWER are adding clumps of pink flowers along streams where moisture is constantly available to the plants.

Shootingstar
NPS/Petrina Vecchio

Marsh marigold, *NPS Photo*

Monkeyflower, *NPS Photo*

Bear grass, NPS Photo

Like the related century plant, BEAR GRASS grows slowly and favors well-drained soils. New bear grass clumps develop from underground stems but they cannot produce flowers for several years until sufficient new starches are produced and reserved in the roots. After the plant blooms once the entire plant dies. Entire communities of bear grass will sometimes go through several summers without blooming – then hundreds will bloom during the same season.

The white flowers and small thickened leaves of TOLMIE SAXIFRAGE form a low-lying mat in rocky areas above timberline. The succulent nature of the leaves allows storage of snow meltwater that can be used later in the summer when the barren rocky outcrops have been drained of moisture.

Alfred Brousseau, Saint May's College

Characteristic of many alpine plants, SPREADING PHLOX forms a dense, low-growing flower cushion which hugs rocks for warmth and wind protection. The smooth, convex surface, like an airfoil, eases wind over its flower cushion with a minimum of resistance. Even so, as wind blows over these mounds it slows, allowing wind-borne particles to fall onto the plant, slowly raising the level of soil within its periphery. At the same time, there is a maximum exposed leaf surface for photosynthesis with minimum exposure to the elements. Temperatures may be several degrees higher inside a cushion plant than outside; the colder it is outside, the greater the difference. Spreading phlox has a deep taproot (8 to 15 feet) which nines water from far below the soil surface.

Phlox, *NPS Photo*

Mat plants, such as PARTRIDGEFOOT, are also specially adapted to subalpine and alpine conditions but are more spreading instead of tightly cushioned. There is little vertical separation between the leaves, thus a happy balance is maintained between the greatest exposure to light and reflected heat from the soil surface and the least exposure to cold, wind damage and drying. Water coming from the roots has the shortest possible distance to travel to the leaf tip. Less of the plant's energy is spent for production of plnt tissue. Dwarfing occurs only in growth shoots; flower size is not affected.

Partridgefoot, *NPS/Steve Redmond*

The white spikes of MOUNTAIN BUSTORT punctuate the subalpine flowerfields of Paradise from spring through autumn. Their "dirty socks" scent carries far on the breeze. Such a distinctive odor may be one way the bistort helps insect pollinators recognize what flower it is.

Bistort, *NPS photo*

Corn lily, *Kelsey Byers*

Heavy beds of snow lying on steep meadows tend to creep down the slope during the winter, scouring off all vegetation at the surface. Woody seedlings are frustrated year after year, while perennial herbs with fat storage roots and fast spring growth, such as the CORN LILY, are favored. Although its flowers are an inconspicuous light green the Corn lily draws a lot of attention with its corn-tassle-like flower heads and large, prominently veined leaves. The young shoots and roots of Corn lily contain toxins which prevent many other species of plants from growing nearby and make it highly poisonous for human consumption. By late summer when the plants mature these protective toxins will drain from the leaves. Deer and insects will then star munching on the plants.

Thirty to forty lavender "petals" typically surround the MOUNTAIN DAYSY's yellow center. The radial symmetry o the flower maximizes the plant's opportunity to absorb sunshine from all directions. This type

Daisy, *NPS photo*

of flower configuration can be distinct advantage in an area with a short growing season.

Fireweed, *Keir Morse*

Fluff balls of silky-winged FIREWEED seeds are carried on the wind to settle in burned areas and avalanche tracks where they will sprout the following summer. Like all plants that have a single flower stalk, the fireweed bloom will progress onward on the stalk so that early-opening flowers will not shade later-opening ones. This blooming pattern increases the amount of sunshine a plant receives during the relatively short growing season typical of high elevations.

Gentian, *NPS/Crow Vecchio*

Last of the flowers to bloom at Paradise, MOUNTAIN BOG GENTIAN grow in small clusters and favor moist meadows and streambanks. The deep blue tubular gentian flower faces upward to the sky and like many gentians it closes when clouds come over or when breathed upon or touched thus protecting its pollen from damage. Pollinating insects are inactive during storms anyway. Since many day insects have an aversion to darkness, tubular flowers like the gentian are often paler toward their bases. Their tiny "windows" make entrance more attractive to pollinating insects.

SEDGE

Black sedge, *Keir Morse*

Late-lying snowbeds allow only an extremely short growing season. Here a plant's ability to grow rapidly is more important than its ability to survive cold. BLACK ALPINE SEDGE is the speed demon among grasslike plants, flowering 5-9 days after its release from the last remnants of the snowpack. Even the quickest seed-setting is rarely quick enough, so it spreads by rhizomes as well. Leaves are reduce to essentials—no bright petals, just the reproductive parts enclosed in a simple protective envelope. Also, the narrow leaves and stems of sedges are not likely to be torn by wind. Being wind pollinated, sedges do not need colorful petals nor nectar nor volatile aromatic oil to attract pollinators, but instead produce pollen in superabundance. For wind pollination to be efficient there must be many individuals of one kind, preferably massed close to each other. Sedges are extremely hardy, thriving in the cooler microclimates of the meadows.

SHOWY SEDGE is the most abundant subalpine tall sedge at Mount Rainier. It grows in meadows with moderate conditions- fairly early snowmelt, and deep soil.

Jewel lichen, *James Lindsey at Ecology of Commanster*

LICHEN

Lichens are formed from a fungus and an alga; fungus forms the

tough outer layers while the inner layers contain algal cells enmeshed in fungal threads. Scholars today question whether the lichen symbiosis is mutually beneficial. Some call it *controlled parasitism* in which the fungi slowly kill their green partners. Lichen, crusted on rocks or clinging to soil, have the ability to absorb more than their own weight of water after a dry spell. They can carry on food production at any temperature above 32 degrees F and can grow even where snowbanks persist very late into the summer. Lichens that colonize bare rock may occasionally, after centuries of life, death, and decomposition, alter the rock enough for plants to grow. But the ability of lichens to break down rock, initiating the long, slow process of succession, has been overrated. It is negligible compared t the effects of frost. Red-orange JEWEL LICHEN marks a pika or marmot perch. The extra nitrate from the animal's urine makes the rock an ideal growing site for this lichen.

The pale, green-grey festoons found on trees from mid-elevations to timberline is known as Old-man's–beard or GOAT'S BEARD LICHEN. Although it is eaten by deer, the apparent browse line at higher elevations is more likely a marker of spring snowpack depth. It relies on spring as a growing season, when day length and temperatures are rising but rain is plentiful.

Goat's beard lichen, *NPS Photo*

Lichen's diet of airborne solids makes them extremely sensitive to airborne sulphur dioxide and other pollutants. No lichens survive in the worst urban air. Environmental scientists watch lichen species as a measure of pollution in an airshed over a period of years.

BIRDS

Succulent saxifrage leaves, white heather flowers and seeds make up the bulk of the summer diet for ROSY FNICHES. They will also forage on glaciers for insects that collapse on the snow, numbed by the snow after being carried astray by upvalley winds. Finches nest in high rock crevices and glacial crevasses. Rosy finches have been seen as high as 11,000 feet on Mount Rainier but are often found during the summer near Glacier Vista (6300 feet) and Panorama Point (6800 feet). Most of our rosy finches winter east of the Cascades. The finch family name comes from the same Latin root as does the word *frigid*.

Rosy finch, *NPS Photo*

Ranging as far north as Alaska, RUFOUS HUMMINGBIRDs are the most northerly of hummingbirds. They are common in subalpine and alpine areas on Mount Rainier from spring through fall and are especially attracted to red flowers such as magenta paintbrush. Ounce for ounce a hummingbird flying has ten times the caloric requirements of a person running. They consume half their body weight in sugar each day. Their needlelike beaks

Male Rufous Hummingbird
Dean E. Biggins/ U.S.F.W.S.

probe into flowers for nectar which they sip for calories. For protein hummingbirds eat small insect and spiders. Rufous hummingbirds, as well as other hummers, become torpid at night to conserve calories. Torpidity is a state in which the body temperature drops to that of the surrounding air, and both the heart rate and respiration rate decrease dramatically. When nighttime temperatures drop below 60 degrees F, the birds may save as much as 98% of the energy they would otherwise use to maintain their normal body temperature. To avoid winter, these hummingbirds will migrate as far as South America.

Ptarmigan, *NPS photo*

Although smaller than their grouse relatives, WHITE-TAILED PTARMIGANs are the largest creatures that make the alpine zone their permanent home. Like other grouse, they rely on camouflage to protect them from predators. In winter ptarmigans molt to pure white plumage and stay on snow as much as possible, crouching low so even their shadow will be smaller and less conspicuous. Ptarmigan's feet and legs are so heavily clad in feathers during winter that they look like pantaloons. The feathers act as snowshoes and provide warm insulation from the cold snow. Once their summer plumage grows in, ptarmigan sty off the snowfields. Adult ptarmigan seem to know when they are changing color and remain on the periphery of snowbanks during spring and fall molts, in a half-and-half environment that matches their neither-nor plumage. By summer their mottled gray plumage allows them to melt into the background of their alpine home. Mating time is triggered by the lengthening daylight of spring. Breeding begins sometime in early may. Ptarmigan are monogamous during the breeding period, with a pair bond usually made for life. Once breeding is completed they have no further interest in each other until the following year, when they find each other again. Weather causes the greatest egg mortality, although nutcrackers, falcons and weasels often find ptarmigan eggs when mother is off the nest. Panorama point is a likely location to look for white-tailed ptarmigan.

Blue grouse, *NPS photo*

During spring and early summer one is bound to hear a constant, deep hooting sound while hiking at Paradise. No, it is not an owl nor a bear as many people imagine, but a male BLUE GROUSE courting a female. Both this courting "hoot" and the chief visual display of males--bare yellow patches on the neck—are performed by inflating a pair of air sacs in the throat. Blue grouse feed and nest on the ground and fly only in infrequent bursts. Grouse relish berries and insects when available during summer but subsist mostly on conifer needles during winter. Like ptarmigan, grouse have feather-insulated feet and appendages which serve as snowshoes to ensure their winter mobility.

VARIED THRUSH are common in the forest and subalpine areas of Mount Rainier. They are milar to robins in appearance but can be identified by their distinctive orange eye stripe and black breast band. Also distinctive to varied thrush is their long and forceful single-note trill.

Varied thrush (NPS/ONP)

Grey jay, NPS

Known by the alias *camp robber*, GRAY JAY has opportunistic feeding habits and can be very aggressive. Loose, fluffy plumage makes the gray jay almost silent in flight. It is abundant in forest and subalpine areas of Mount Rainier and can be seen at Paradise throughout most of the year.

Clark's nutcracker, NPS

Although sometimes mistaken for a gray jay, CLARK'S NUTCRACKER is a much larger bird with a long, straight, black beak used to crack seeds from coniferous trees.

Dark-eyed junco/Will Elder, NPS

DARK-EYED JUNCOs are abundant in our forests and subalpine areas. The white outer tail feathers of this six-inch resident bird flash when the junco is in flight.

Raven, NPS

Western expansion pushed the COMMON RAVEN up to the mountains from the lowlands where it was once as abundant as its relative the crow. Ravens share the opportunistic feeding habits of other jays but also act like raptors consuming rodents, rabbits, insects, worms, nesting birds, moles and frogs.

MAMMALS

Easily identified by the stripes along their cheeks as well as along the sides of their bodies, chipmunks are the smallest of the pacific Northwest squirrels. It is the YELLOW PINE CHIPMUNK which inhabits the subalpine areas. Chipmunks are characteristically active during daylight hours and spend most of their time on the ground or in low bushes. They are not often found in the open habitats preferred by ground squirrels and chipmunks seldom climb trees. They feed on seeds, leaves, flower stems and fruit. Chipmunks do not typically depend on a deposit of fat to supply energy during winter. Instead they begin storing food in their dens starting late September. Their cache of seeds, leaves, flower

Chipmunk/Canter 1972

stems, and fruit is covered over with a layer of grass for better storage. Although chipmunks hibernate, from time to time during the winter they dig down through the grass and consume some of their stores.

Golden-mantled ground squirrel, NPS

GOLDEN-MANTLED GROUND SQUIRREL can be identified by the narrow white line above and below the eye and stripes along its sides. This little squirrel is very much like a chipmunk in habits, behavior and ecology.

Of the five species of marmots represented in the Pacific Northwest, it is the HOARY MARMOT which inhabits subalpine areas. A marmot is active for oly a few months of the year. It spends its time sunning on rocks and eating flowers and stems, grasses and succulent herbs. The ody fat it adds to its back and tail during summer allows the marmot to spend the winter in hibernation. Hibernation can be thought of as a form of seasonal migration. The primary impetus for hibernation is probably lowering air temperature. During hibernation, a marmot's heart beats only four to five times a minute. One full breath is taken about every minute. Body temperature is only a few degrees above freezing. Marmots inhabit rocky places where they find shelter in rock crevices and underground burrows. Eagles, cougars, bobcats, coyotes and bears are their enemies. Since they have no way to defend themselves but to run for their burrows, when danger is sighted a marmot will issue a shrill whistle to warn its compatriots. Hoary marmots are the largest members of the squirrel family.

Hoary marmot, NPS

PIKAs, small members of the Rabbit family, are about the size of guinea pigs. They inhabit talus slopes and rocky outcrops between 3000 feet and 8000 feet. The soles of their feet are densely furred to provide traction for scampering across slippery rocks. Pikas venture into nearby meadows only a short ways to gather leaves and flower stems which are carried back to the rocks and stacked in small piles for drying in the sun. Pikas dry their food slowly and carefully so it becomes completely dry before being stored underground. They even take care to move their food piles underground when rain threatens. This prevents their food from rotting during winter. A pika's vegetarian diet is not high in calories. The animal must fill its stomach almost hourly to meet its energy needs. Pikas do not hibernate during the winter as chipmunks, ground squirrels and marmots do, instead they stay active under the snow among the rocks. Temperatures beneath the snow cover remain around 26 degrees F, much warmer than the outside air temperature might be during most of the winter. Their ears are small and rounded, less likely to freeze. Their tail is so small as to be non-existent, a common characteristic among winter-active animals since small extremities help conserve body heat.

Pika, NPS

SNOWSHOE HARE is also active throughout the winter but, unlike the pika who is cozy among the rocks under the snow, the hare is active on top of the snow. It doesn't even build a nest or use a burrow to sty warm but retires to shallow depressions called *forms* under shrubs and low tree branches. The diet of the snowshoe hare changes from one of fresh green s during summer to one of conifer buds and shrub bark during winter. The shortening length of the autumn days triggers an internal clock in the snowshoe hare. The hare begins to molt beginning with the tips of its ears and its hind feet. During the autumn molt, its lightweight brown summer fur is replaced by a heavy coat of white winter fur, providing both extra protection against cold temperatures and protective camouflage to confuse predators. Stiff hairs grow between its toes and along the edges of its feet serving the animal as snowshoes. The molt may take weeks to complete but by the time the snow covers the ground, the snowshoe hare is completely white. Come spring, the lengthening days will trigger a return to summer pelage. Snowshoe hares can attain speeds up to 25 miles per hour over short distances and can leap up to 10 feet in a single bound.

Snowshoe hare, NPS

BLACK BEAR is the smallest and he most common of the North American bears. Although plant foods dominate its diet, bears also eat small mammals, insects or larvae, blueberry fruit and leaves. Bears prepare their dens with fluffy stuff—ideally cedar bark which stays resilient under the prolonged weight of the bear's body. Between the insulative nest and superlatively thick fur, the bear maintains a body temperature of about 88 degrees F throughout winter, however heart weight may reach a low of eight beats per minute. Typically two or three cubs are born around January. The mother wakes up to give birth, then nurses them mostly in her sleep for the next few months. Even though black bears normally withdraw from any contact with people, bears should always be considered dangerous—especially sows with cubs. Black bears are good tree climbers, swim well and can run 25 mph for short stretches.

Black bear, NPS

BLACKTAILED DEER and mule deer were considered two distinct species until fairly recently. Now they are lumped together. In winter deer seek thermal cover on steep south aspects just above river bottoms. The low elevation and insulating conifer canopy offer warmer temperatures, shallower snow, and more shrubs to browse. As snow melts, deer move up slope to subalpine meadows. Antlers, made of solid bone and usually branched, are a defining characteristic of the deer family. In late winter the antlers weaken at their bases and fall off. A new pair begins to grow by early summer. Antlers exist to help establish dominance among males for access to females during the rut.

Blacktailed deer, NPS

RED FOXes prefer timberline and other areas of broken forest cover. They are largely nocturnal, shy and elusive but re often seen at Paradise during winter nights by campers who have left food outside their tents and igloos. An adult fox does not sleep in a dean during winter. Instead a fox will curl its tail around

its nose to form a warm furry package. The fox may become covered by snow during a storm—perhaps some of the small, snowy hummocks skiers see during winter re actually sleeping foxes! Reds foxes mate in mid-winter and bear their litter in early spring. Like human siblings, red fox littermates may be blond, dark or red. All have white tail tips. Foxes prefer to eat mice, hares, frogs, squirrels and young marmots but also eat lots of insects, fruit and seeds.

Red fox, NPS

REFERENCES

ANIMAL TRACKS OF THE PACIFIC NORTHWEST
Karen Pandell, Chris Stall

BIRDS OF THE PACIFIC NORTHWEST MOUNTAINS
Jan L. Wassink

CASCADE-OLYMPIC NATURAL HISTORY
Daniel Mathews

FORESTS OF MOUNT RAINIER
William H. Moir

LAND ABOVE THE TREES
Ann Zwinger, Beatrice Wilard

MAMMALS OF THE NORTHWEST
Earl J. Larrison

MOUNTAIN FLOWERS
Harvey Manning

NORTHWEST MOUNTAIN WEATHER
Jeff Renner

TIMBERLINE
Stephen F. Arno, Ramona P. Hammerly

WILDFLOWERS OF THE PACIFIC NORTHWEST
Lewis J. Clark

KEY REFERENCES, PACIFIC NW ALPINE & SUBALPINE VEGETATION

Belsky, J., and del Moral, R. 1982. Ecology of an alpine-subalpine meadow complex in the Olympic Mountains, Washington. *Canadian Journal of Botany*. 60: 779-788

Bliss, L.C. 1969. Alpine community patterns in relation to environmental parameters *In* Greenridge, K.N.H. (ed.), *Essays in Plant Geography and Ecology*. Helikon. Nova Scotia Museum, 167-184.

Canaday, B.B. and R.W. Fonda. 1974. The influence of subalpine snowbanks on vegetation pattern, production, and phenology. *Bulletin of the Torrey Botanical Club* 101: 240-350. [Olympic Mountains]

del Moral, R. 1979a. High elevation vegetation of the Enchantment Lakes Basin, Washington. *Canadian Journal of Botany* 57: 1117-1120.

_____. 1979b. High elevation species clusters in the Enchantment Lakes Basin. *Madroño* 26: 164-172.

_____. 1983a. Competition as a control mechanism in subalpine meadows. *American Journal of Botany* 70: 232-245. [Olympic Mountains]

_____. 1983b. Vegetation colonization of subalpine meadows using adaptive strategies. *Canadian Journal of Botany* 61: 3117-3127. [Olympic Mountains]

Douglas, G.W. 1972. Subalpine plant communities of the western North Cascades, Washington. *Arctic and Alpine Research* 4: 147-156.

_____ and L.C. Bliss. 1977. Alpine and high subalpine plant communities of the North Cascades Range, Washington and British Columbia. *Ecological Monographs* 47: 113-150.

Edwards, O.M. 1980. *The Alpine Vegetation of Mount Rainier National Park: Structure, Development, and Constraints*. Ph.D. dissertation. University of Washington, Seattle. 264 pp.

Fonda R.W. 1975. Ecology of alpine timberline in Olympic National Park. *Proceedings of the Conference for Scientific Research in the National Parks* 1: 209-212.

_____ and L.C. Bliss. 1969. Forest vegetation of the montane and subalpine zones, Olympic Mountains, Washington. *Ecological Monographs* 39: 271-301.

Franklin, J.F., and C.T. Dyrness. 1988. *Natural Vegetation of Oregon and Washington*. Oregon State University Press. 452 pp. (Includes many ref. for Mt. Rainier)

Kuramoto, R.T., and Bliss, L.C. 1970. Ecology of subalpine meadows in the Olympic Mountains, Washington. *Ecological Monographs*, 40: 317-347.

Pfitsch, W.A., and L.C. Bliss. 1985. Seasonal forage availability and potential vegetation limitations to a mountain goat population, Olympic National Park. *American Midland Naturalist* 113: 109-121.

Schreiner, E.G. 1994. Subalpine and alpine plant communities. In D.B. Houston, E.G. Schreiner, and B.M. Moorhead, eds., *Mountain Goats in Olympic National Park: Biology and Management of an Introduced Species*. Scientific Monograph NPS/NROLYM/NRSM-94/25, U.S. Department of Dedistorior, National Park Service. 242-250.

Taylor, R.J., and G.W. Douglas. 1978. Plant ecology and natural history of Chowder Ridge, Mt. Baker: a potential alpine research natural area in the western North Cascades. *Northwest Science* 52: 35-40.