

# Buffelgrass Facts and Figures

January, 2018

## Threats to the Ecosystem

**Fire** is the **greatest threat** because of radical changes in fire regime in desert areas.

- Buffelgrass will allow **more frequent fires**.<sup>47,48</sup>
  - Buffelgrass creates **combustible fuel** that is present year round (it is a perennial plant).
    - The Sonoran Desert historically burned every ~ 250 years or more when annual plants boomed after wet periods, and even then fires were mild and patchy.<sup>1-4</sup>
    - **Frequency will increase** because of continuous grass that allows fire spread from backyards to natural areas.<sup>33,35-37,47,48</sup>
- Fire **size will increase** because fuel allows fire to spread further, burning more area.<sup>2, 5-7, 47,48</sup>
- More grass means **increased fire intensity**.
  - Buffelgrass **fuel loads of 1- 4 tons per acre** were found on Saguaro NP.<sup>8,9,62</sup> Similar fuel loads have been recorded in Australia.<sup>59</sup>
    - This is more than **2 times to 4,000 times** usual Sonoran Desert fuel loads (including non-native grasses such as red brome and Mediterranean grass).<sup>8-12</sup>
  - Buffelgrass fire **temperatures** were recorded at **1300-1600 F**<sup>8,62</sup> versus 190 – 750 F recorded in wildfires fueled by desert annual plants.<sup>13</sup>
  - Rate of buffelgrass fire spread was recorded at **2-3 mph** under moderate weather conditions.<sup>8, 61</sup>
    - Predicted rate of spread is 4-48 times faster than usual Sonoran Desert fires.<sup>14</sup>
  - Buffelgrass **flame length recorded at 12-18 ft** under moderate conditions.<sup>8,61</sup>
  - Buffelgrass fires miss very little: 99% of available fuel was burned in one experiment.<sup>8</sup>
- **Damage from wildfires will be unprecedented.**
  - Previous fires in the Sonoran Desert were fueled primarily by non-native annual grasses (red brome/Mediterranean grass),<sup>3,15,16</sup> and resulted in 20-80% saguaro mortality and significant mortality to other desert vegetation including barrel cacti, cholla, and paloverde tree.<sup>1,2,6,10,17-22</sup>
  - Damage from buffelgrass fires will increase to levels not experienced before because of more fuel.<sup>6,9,18,20,23,47</sup>
- Fire damage will likely lead to **increases in other invasive species and buffelgrass**.<sup>31,33,47</sup>

**Signature and common species** of the Sonoran Desert are threatened

- Buffelgrass increases exponentially.<sup>45</sup>
- Untreated areas were documented with a 350% increase of buffelgrass over two years in one place in the park.<sup>50</sup>
- Buffelgrass *directly* affects plants through **competition and habitat alteration**.<sup>6,7,16,24-26, 45,47,48,60</sup>
  - Above 40% buffelgrass cover, native plant cover is very low.<sup>8, 45,61</sup>
  - Large patches of buffelgrass “may be largely devoid of native vegetation.”<sup>45</sup>
- Buffelgrass may have chemical compounds that inhibit germination and growth of other plants.<sup>46,49</sup>
- Buffelgrass *indirectly* affects plants and animals from **fire-induced mortality**.
  - **Native species did not evolve with fire**; many desert plants are fire-intolerant, and animals move too slowly to escape or are not adapted to the fire-altered habitat.<sup>6,22,27-31,47,48</sup>
- Buffelgrass has been documented as reducing condition of desert tortoises through habitat changes, with the potential for more severe long-term impacts.<sup>58</sup>
- Recurrent fires could cause **localized extinction** and result in vegetative type **conversion into a grassland or scrubland**<sup>7,8,9,32-34,61</sup> and an overall **decrease in Sonoran Desert biodiversity**.<sup>16,24,27,28,38,61</sup>

## Public safety and the economy

- **Public safety and private property will be threatened** by intense wildland urban interface fires.<sup>8,14,15,39,62</sup>
- Tucson's **economy will be harmed**, because it depends on tourism, which depends on the Sonoran Desert's unique plants and animals.<sup>44,48</sup>
- **Real estate values may decline** because of the threat of fire and damaged environment.<sup>44</sup>

## Saguaro National Park's Buffelgrass Efforts

Saguaro National Park is among the leaders in the efforts to control buffelgrass.

- Park employees and volunteers are **using herbicides** and **manually pulling and digging up** buffelgrass.
  - Pulling and spraying both require **at least 3 years of repeated treatments** because of seeds left in the soil<sup>41,42</sup>
- We are also **conducting or funding research and monitoring** to find the best ways to control buffelgrass and protect native species.
- Park research shows that herbicide can still effectively kill buffelgrass when the plant is in early dormancy (~50% green).
- The Park uses glyphosate as an herbicide, which must be absorbed through green leaves. The Park has experimented with other herbicides, and has funded research using grass-specific herbicides, but glyphosate still remains the best option currently available—least toxic and most effective.
  - Glyphosate has low toxicity, but it is an eye irritant.<sup>57</sup>
  - The NPS follows EPA guidance on herbicide use. The EPA currently does not consider glyphosate a carcinogen.
  - Results of numerous studies of its links to cancer have been contradictory. The World Health Organization considers it a probable carcinogen<sup>51</sup>, and the state of California do consider it a known carcinogen.<sup>52</sup> WHO places glyphosate in the same class as eating fried food and working a night shift.<sup>53</sup> California places it in the same category as tobacco smoke<sup>52</sup>.
  - The WHO states that the probability of developing a cancer “will depend on factors such as the type and extent of exposure and the strength of the effect of the agent.”<sup>51</sup>
  - Cancer Assessment Review Committee of the EPA found that glyphosate should be labeled as “not likely to be carcinogenic to humans”<sup>54</sup> and the European Food Safety Authority and European Chemicals Agency both determined that glyphosate was unlikely to pose a carcinogenic hazard to humans.<sup>55,56</sup>

### *Facts and figures about Saguaro National Park's efforts:*

- Buffelgrass was introduced into southern Arizona beginning in the 1930's for erosion control and for livestock forage.
- Numerous strains have been collected throughout its range (stretching from Africa to India) in attempts to find species and cultivars to improve forage production and soil conservation. The strain in Southern Arizona is cultivar T-4464. That strain was collected in the Turkana Desert in Kenya in the 1940's by a scientist from South Africa. Seed was shipped to the US Department of Agriculture in 1946. T-4464 was released by the Soil Conservation Service in 1949, and by 1985, seed companies in Texas had sold almost 8,000 tons of seed, and Texas ranchers had established it on about 10 million acres of land.<sup>63, 64</sup>
- 1970's-1980's, buffelgrass widely planted in the area for erosion control.
- **1989 first known observation** of buffelgrass in the Park (photograph from saguaro monitoring)
- 1991 first written record of buffelgrass in the Park
- 1993 staff and volunteers **begin manual removal of buffelgrass**

- 2000 increased awareness and park attention to buffelgrass; establishment of Invasive Plant Program
- 2002 extensive survey of buffelgrass at both districts estimated ~175 acres of buffelgrass
- 2003 spatial logistic regression model produced which predicts buffelgrass could occupy up to 100% of the Arizona Upland of the Sonoran Desert biome within the park <sup>42</sup>
- 2002-2004 research project to evaluate various control methods
- 2004 Environmental Assessment and Exotic Plant Management Plan
- **2005 begin use of herbicides to control buffelgrass**, complementing manual removal
- Fall 2012, estimate of **over 2,000 acres of buffelgrass** in the park
- 2013-2014 Environmental Assessment used to evaluate potential use of aerial spraying for areas too remote or too rugged for ground treatments
- **2014 aerial spraying of buffelgrass began**, complementing ground spraying and manual removal.
  - Numerous mitigations enacted to protect human safety and natural and cultural resources.
    - No-spray zones around private property (1/8 mile from private land and 1/4 mile from occupied buildings).
    - Large droplet size to prevent drift, keeping the herbicide on target.
    - Strict weather guidelines, especially wind, to reduce drift.
    - Spraying where buffelgrass is 50% or more of the plant cover. (Few native plants are present in these dense patches.)
  - Three years of intense monitoring has found that aerial spraying reduces buffelgrass dominance, and that saguaros and palo verdes suffered minimal damage.<sup>50</sup> This is probably due to the waxy cuticle present on cacti and small leaves and differences in metabolism in palo verde.
  - 2017 will be the final year of intense monitoring, but long-term monitoring of saguaros will continue
- Volunteers contribute over 3,000 hours annually to help control buffelgrass
- Estimated costs average \$500/acre (range \$200-\$600/acre) for chemical treatments

## Community Efforts

The fight against buffelgrass is a **community-wide** effort.

- Multi-partner Buffelgrass Working Group implementing the Southern Arizona Buffelgrass Strategic Plan
- Pima County hosts educational trainings and speakers bureau; developed outreach materials
  - See: <https://tucsoncleanandbeautiful.org/buffelgrass-education-removal/>
- Pima County ordinance to control buffelgrass when deemed a threat to human safety and property
- Rural Metro developing strategies for prevention/education/mitigation, and modifying training to reflect new fuel source
- Region-wide volunteer efforts to map and pull buffelgrass, and numerous “weed wacking” groups in the area (see <http://www.desertmuseum.org/buffelgrass/volunteer.php>)
- Annual, community-wide Beat Back Buffelgrass Day, with over 1,000 volunteers per year.

## Contacts at Saguaro National Park

Perry Grissom, Restoration Ecologist

Phone: 520-733-5179

Email: [perry\\_grissom@nps.gov](mailto:perry_grissom@nps.gov)

## Resources

[www.buffelgrass.org](http://www.buffelgrass.org)

<http://www.nps.gov/sagu/naturescience/invasive-plants.htm>

## References

- <sup>1</sup> Humphrey, R.R. 1974. Fire in the deserts and desert grassland of North America. Pages 366-400, in *Fire and Ecosystems* (eds. T.T. Kozlowski and C.E. Ahlgren). Academic Press, New York.
- <sup>2</sup> McLaughlin, S.P. and J.E. Bowers. 1982. Effects of wildfire on a Sonoran Desert plant community. *Ecology* 63:246-248.
- <sup>3</sup> Schmid, M.K. and G.F. Rogers. 1988. Trends in fire occurrence in the Arizona Upland subdivision of the Sonoran Desert, 1955 to 1983. *Southwestern Naturalist* 33:437-444.
- <sup>4</sup> Schussman, H., C. Enquist, and M. List. 2006. Historic fire return intervals for Arizona and New Mexico: a regional perspective for southwestern land managers. *The Nature Conservancy in Arizona*.  
[http://azconservation.org/downloads/data/historical\\_fire\\_return\\_intervals\\_for\\_arizona\\_and\\_new\\_mexico](http://azconservation.org/downloads/data/historical_fire_return_intervals_for_arizona_and_new_mexico)
- <sup>5</sup> Wright, H.A. and A.W. Bailey. 1982. *Fire ecology: United States and southern Canada*. Wiley and Sons, New York.
- <sup>6</sup> Búrquez-Montijo, A., M.E. Miller, and A. Martínez-Yrizar. 2002. Mexican grasslands, thornscrub, and transformation of the Sonoran Desert by invasive exotic buffelgrass (*Pennisetum ciliare*). Pages 126-146 In Tellman, B., editor, *Invasive exotic species in the Sonoran Region*. University of Arizona Press and The Arizona-Sonora Desert Museum, Tucson, AZ.
- <sup>7</sup> Clarke, P.J., P.K. Latz, and D.E. Albrecht. 2005. Long-term changes in semi-arid vegetation: invasion of an exotic perennial grass has larger effects than rainfall variability. *Journal of Vegetation Science* 16:237-248.
- <sup>8</sup> McDonald, C. 2009. Management of non-native perennial grass in southern Arizona: effects of prescribed fire and livestock grazing. PhD Dissertation, University of Arizona.
- <sup>9</sup> Esque, T., C. Schwalbe, J.A. Lissow, D.F. Haines, D. Foster, and M.C. Garnett. 2007. Buffelgrass fuel loads in Saguaro National Park, Arizona, increase fire danger and threaten native species. *Park Science* 24:33-56.
- <sup>10</sup> Cave, G.H. and D.T. Patten. 1984. Short-term vegetation responses to fire in the Upper Sonoran Desert. *Journal of Range Management* 37:491-496.
- <sup>11</sup> Halvorson, W.L. and D.T. Patten. 1975. Productivity and flowering of winter ephemerals in relation to Sonoran Desert shrubs. *American Midland Naturalist* 93:311-319.
- <sup>12</sup> Patten, D.T. 1978. Productivity and production efficiency of an Upper Sonoran Desert ephemeral community. *American Journal of Botany* 65:891-895.
- <sup>13</sup> Patten, D.T. and G.H. Cave. 1984. Fire temperatures and physical characteristics of a controlled burn in the Upper Sonoran Desert. *Journal of Range Management* 37:277-280.
- <sup>14</sup> Estimated with conditions representing average June weather for Tucson using fire behavior prediction program BehavePlus4.0 (Andrews, Patricia L.; Bevins, Collin D.; Seli, Robert C. 2008. BehavePlus fire modeling system, version 4.0: User's Guide. Gen. Tech. Rep. RMRS-GTR-106WWW Revised. Ogden, UT:

Department of Agriculture, Forest Service, Rocky Mountain Research Station). Buffelgrass modeled with fuel model GR6, Sonoran Desert vegetation modeled with fuel models GR1, GS1, GS2, SH1, and SH2 (Scott, J.H. and R.E. Burgan. 2005). Standard fire behavior fuel models: a comprehensive set for use with Rothermel's surface fire spread model. USDA Forest Service Rocky Mountain Research Station General Technical Report RMRS-GTR-153.) with inputs (1hr = 4, 10hr = 5, 100hr = 6, live herbaceous = 30, live woody = 50, mid-flame windspeed 0, 3, 6, 9, 12, and 15 mph, slope = 0). Modeling conducted 1/6/10 by Perry Grissom, Fire Ecologist, Saguaro NP.

<sup>15</sup> Van Devender, T.R., R.S. Felger, and A. Búrquez M. 1997. Exotic plants in the Sonoran Desert Region, Arizona and Sonora. California Exotic Pest Plant Council, 1997 Symposium Proceedings.

<sup>16</sup> Williams, D.G., and Z. Baruch. 2000. African grass invasion in the Americas: ecosystem consequences and the role of ecophysiology. *Biological Invasions* 2:123-140.

<sup>17</sup> Alford, E.J., J.H. Brock, and G.J. Gottfried. 2005. Effects of fire on Sonoran Desert plant communities. Pages 451-454 in Gottfried, G.J., B.S. Gebow, L.G. Eskew, and C.B. Edminster (compilers). *Connecting mountain islands and desert seas: biodiversity and management of the Madrean Archipelago II*. Fifth Conference on research and resource management in the southwestern deserts. RMRS-P-36. USDA Forest Service, Ft. Collins, CO.

<sup>18</sup> Esque, T.C., C.R. Schwalbe, D.F. Haines, and W.L. Halvorson. 2004. Saguars under siege: invasive species and fire. *Desert Plants* 20:49-55.

<sup>19</sup> Reynolds, H.G. and J.W. Bohning. 1956. Effects of burning on a desert grass-shrub range in southern Arizona. *Ecology* 37:769-777.

<sup>20</sup> Rogers, G.F. 1985. Mortality of burned *Cereus giganteus*. *Ecology* 66:630-632.

<sup>21</sup> Rogers, G.F. and J. Steele. 1980. Sonoran Desert fire ecology. Pages 15-19 In M.A. Stokes and J.H. Dieterich (editors), *Proceedings of the fire history workshop*. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station General Technical Report RM-GTR-81, Fort Collins, CO.

<sup>22</sup> Wilson, R.C., M.G. Narog, A.L. Koonce, and B.M. Corcoran. 1995. Postfire regeneration in Arizona's giant saguaro shrub community. Pages 424-431 in L.F. DeBano, G.J. Gottfried, R.H. Hamre, C.B. Edminster, P.F. Ffolliott, and A. Ortega-Rubio, Technical Coordinators. *Proceedings of biodiversity and management of the Madrean Archipelago*. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. General Technical Report RM-GTR-264.

<sup>23</sup> Thomas, P.A.. 1991. Response of succulents to fire: a review. *International Journal of Wildland Fire* 1:1-22.

<sup>24</sup> Burgess, T.L., J.E. Bowers, and R.M. Turner. 1991. Exotic plants at the desert laboratory, Tucson, Arizona. *Madroño* 38:96-114.

<sup>25</sup> De La Barrera, E. and A.E. Castellanos. 2007. High temperature effects on gas exchange for the invasive buffel grass (*Pennisetum ciliare* [L.] Link). *Weed Biology and Management* 7:128-131.

<sup>26</sup> Morales-Romero, D. and F. Molina-Freaner. 2008. Influence of buffelgrass pasture conversion on the regeneration and reproduction of the columnar cactus, *Pachycereus pectin-aboriginum*, in northwestern Mexico. *Journal of Arid Environments* 72:228-237.

- <sup>27</sup> Bestelmeyer, B.T. and R.L. Schooley. 1999. The ants of the southern Sonoran desert: community structure and the role of trees. *Biodiversity and conservation* 8:643-657.
- <sup>28</sup> Brooks, M.L. and T.C. Esque. 2002. Alien plants and fire in desert tortoise (*Gopherus agassizii*) habitat of the Mojave and Colorado Deserts. *Chelonian Conservation and Biology* 4:330-340.
- <sup>29</sup> Esque, T.C., C.R. Schwalbe, L.A. DeFalco, R.B. Duncan, and T.J. Hughes. 2003. Effects of desert wildfires on desert tortoise (*Gopherus agassizii*) and other small vertebrates. *Southwestern Naturalist* 48:103-111.
- <sup>30</sup> Flanders, A.A., W.P. Kuvlesky, Jr., D.C. Ruthven III, R.E. Zaiglin, R.L. Bingham, T.E. Fulbright, F. Hernández, and L.A. Brennan. 2006. Effects of invasive exotic grasses on South Texas rangeland breeding birds. *The Auk* 123:171-182.
- <sup>31</sup> Rice, P.M., G.R. McPherson, and L.J. Rew. 2008. Fire and nonnative invasive plants in the Interior West Bioregion. Pages 141-173 In Zouhar, K., J.K. Smith, S. Sutherland, and M.L. Brooks. *Wildland fire in ecosystems: fire and nonnative invasive plants*. Gen. Tech. Rep. RMRS-GTR-42-vol. 6. USDA, Forest Service, Rocky Mt. Res. Sta., Ogden, UT.
- <sup>32</sup> Brooks, M.L., C.M. D'Antonio, D.M. Richardson, J.B. Grace, J.E. Kelley, J.M. DiTomaso, R.J. Hobbs, M. Pellant, and D. Pyke. 2004. Effects of invasive alien plants on fire regimes. *BioScience* 54:677-688.
- <sup>33</sup> D'Antonio, C.M. and P.M. Vitousek. 1992. Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics* 23:63-87.
- <sup>34</sup> Friefelder, R.R., P.M. Vitousek, and C.M. D'Antonio. 1998. Microclimate change and effect on fire following forest-grass conversion in seasonally dry tropical woodland. *Biotropica* 30:286-297.
- <sup>35</sup> Cardille, J.A., S.J. Ventura, and M.G. Turner. 2001. Environmental and social factors influencing wildfires in the Upper Midwest, United States, *Ecological Applications* 11:111-127.
- <sup>36</sup> Radeloff, V.C., R.B. Hammer, S.I. Stewart, J.S. Fried, S.S. Holcomb, and J.F. McKeefry. 2005. The wildland-urban interface in the United States. *Ecological Applications* 15:799-805.
- <sup>37</sup> Syphard, A.D., V.C. Radeloff, J.E. Keeley, T.J. Hawbaker, M.K. Clayton, S.I. Stewart, and R.B. Hammer. Human influence on California fire regimes. *Ecological Applications* 17:1388-1402.
- <sup>38</sup> Franklin, K.A., K. Lyons, P.L. Nagler, D. Lampkin, E.P. Glenn, F. Molina-Freaner, T. Markow, and A.R. Huete. 2006. Buffelgrass (*Pennisetum ciliare*) land conversion and productivity in the plains of Sonora, Mexico. *Biological Conservation* 127:62-71.
- <sup>39</sup> Brooks, M.L. 2008. Plant invasions and fire regimes. Pages 33-45 In Zouhar, K., J.K. Smith, S. Sutherland, and M.L. Brooks. *Wildland fire in ecosystems: fire and nonnative invasive plants*. Gen. Tech. Rep. RMRS-GTR-42-vol. 6. USDA, Forest Service, Rocky Mt. Res. Sta., Ogden, UT.
- <sup>40</sup> Rogstad, A. (editor) 2008. Southern Arizona buffelgrass strategic plan. Prepared by the Buffelgrass Working Group. Tucson, AZ. Available from: <http://www.buffelgrass.org/strategicplan.php>.

- <sup>41</sup> Hacker, J.B. The potential for buffel grass renewal from seed in 16-year-old buffel grass-siratro pastures in south-east Queensland. *Journal of Applied Ecology* 26:213-222.
- <sup>42</sup> Winkworth, R.E. Longevity of buffel grass seed sown in an arid Australian range. *Journal of Range Management* 24:141-145.
- <sup>43</sup> Ward, J.P. 2003. Estimating the potential distribution of buffelgrass in Saguaro National Park, Arizona: Illustration of conservation planning tool in the age of biotic homogenization. M.S. University of Arizona, Tucson. 86pp.
- <sup>44</sup> Buffelgrass Invasion: Its Threat to the Sonoran Desert and Our Will to Stop It. 2009. Video at <https://www.youtube.com/watch?v=nQtIVzSrqZY>. Produced by Mark Betancourt.
- <sup>45</sup> Olsson, A.D., J. Betancourt, M.P. McClaran, and S.E. Marsh. 2012. Sonoran Desert ecosystem transformation by a C<sub>4</sub> grass without the grass/fire cycle. *Diversity and Distributions* 18:10-21.
- <sup>46</sup> Hussain, F., I. Ilahi, S. A. Malik, A.A. Dasti, and B. Ahmad. 2011. Allelopathic effects of rain leachates and root exudates of *Cenchrus ciliaris* L. and *Bothriochloa pertusa* (L.) A. Camus. *Pakistan Journal of Botany* 1: 341-350.\*
- <sup>47</sup> Marshall, V.M., M.M. Lewis, and B. Ostendorf. 2012. Buffel grass (*Cenchrus ciliaris*) as an invader and threat to biodiversity in arid environments: A review. *Journal of Arid Environments* 78:1-12.
- <sup>48</sup> Stevens, J., and D.A. Falk. 2008. Can buffelgrass invasions be controlled in the American Southwest? Using Invasion Ecology Theory to understand buffelgrass success and develop comprehensive restoration and management. *Ecological Restoration* 27:417-427.
- <sup>49</sup> Fulbright, N. and T.E. Fulbright. 1990. Germination of 2 legumes in leachate from introduced grasses. *Journal of Range Management* 43:466-467.
- <sup>50</sup> Jonas, J.L. 2017. Draft report: Aerial application of herbicide for buffelgrass (*Cenchrus ciliaris*) control at Saguaro National Park: Year 3 (2016) monitoring results. 37 pages.
- <sup>51</sup> International Agency for Research on Cancer (World Health Organization). 2016. Q&A on Glyphosate. [https://www.iarc.fr/en/media-centre/iarcnews/pdf/Q&A\\_Glyphosate.pdf](https://www.iarc.fr/en/media-centre/iarcnews/pdf/Q&A_Glyphosate.pdf).
- <sup>53</sup> University of Michigan, Risk Science Center. 2015. Video: “What does ‘Probably Cause Cancer’ actually mean?” <https://www.youtube.com/watch?v=CbBkB81ySxQ>
- <sup>54</sup> Environmental Protection Agency. 2016. Glyphosate Issue Paper: Evaluation of Carcinogenic Potential (glyphosate). 227 pages. [https://www.epa.gov/sites/production/files/2016-09/documents/glyphosate\\_issue\\_paper\\_evaluation\\_of\\_carcinogenic\\_potential.pdf](https://www.epa.gov/sites/production/files/2016-09/documents/glyphosate_issue_paper_evaluation_of_carcinogenic_potential.pdf)
- <sup>55</sup> European Food Safety Authority. 2015. EFSA explains risk assessment: glyphosate. [http://www.efsa.europa.eu/sites/default/files/corporate\\_publications/files/efsaexplainsglyphosate151112en.pdf](http://www.efsa.europa.eu/sites/default/files/corporate_publications/files/efsaexplainsglyphosate151112en.pdf). 4 pages.
- <sup>56</sup> European Chemicals Agency. 2017. Glyphosate not classified as carcinogen by ECHA. <https://echa.europa.eu/-/glyphosate-not-classified-as-a-carcinogen-by-echa>.

- <sup>57</sup> Henderson, A.M., J.A. Gervais, B. Luukinen, K. Buhl, and D. Stone. 2010. Glyphosate General Fact Sheet. National Pesticide Information Center, Oregon State University Extension Services. <http://npic.orst.edu/factsheets/glyphogen.html>. Phone 1-800-858-7378.
- <sup>58</sup> Gray, K.M., and R.J. Steidl. 2015. A plant invasion affects condition but not density or population structure of a vulnerable reptile. *Biological Invasions* 17:1979-1988.
- <sup>59</sup> McKenna, P., V. Glenn, P.D. Erskine, D. Doley, and A. Sturgess. 2017. Fire behavior on engineered landforms stabilized with high biomass buffel grass. *Ecological Engineering* 101:237-246.
- <sup>60</sup> Bracamonte, J.A., C. Tinoco-Ojanguren, M.E. Sanchez Coronado, and F. Molina-Freaner. 2017. Germination requirements and the influence of buffelgrass invasion on a population of *Mammillaria grahamii* in the Sonoran Desert. *Journal of Arid Environments* 137:50-59.
- <sup>61</sup> McDonald, C.J., and G.R. McPherson. 2011. Fire behavior characteristics of buffelgrass-fueled fires and native plant community composition in invaded patches. *Journal of Arid Environments* 75:1147-1154.
- <sup>62</sup> McDonald, C.J., and G.R. McPherson. 2013. Creating hotter fires in the Sonoran Desert: buffelgrass produces copious fuels and high fire temperatures. *Fire Ecology* 9:26-39.
- <sup>63</sup> Cox, J.R., M.H. Martin-R, F.A. Ibarra-F, J.H. Fourie, J.F.G. Rethman, and D.G. Wilcox. 1988. The influence of climate and soils on the distribution of four African grasses. *Journal of Range Management* 41:127-139.
- <sup>64</sup> Ibarra-F., F.A., J.R. Cox, M.H. Martin-R., T.A. Crowl, and C.A. Call. 1995. Predicting buffelgrass survival across a geographical and environmental gradient. *Journal of Range Management* 48:53-59.

\* Note: There is disagreement in the taxonomy of buffelgrass. In Asia and Australia and in older American literature, buffelgrass is called *Cenchrus ciliaris*, but most American literature calls it *Pennisetum ciliare*.