



Abraham Lincoln Birthplace National Historical Park

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Abraham Lincoln Birthplace National Historical Park (hereafter, the Park) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Park, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1). Among the species likely to be found at the Park today, climate suitability in summer under the high-emissions pathway is projected to improve for 8, remain stable for 17, and worsen for 6 species. Suitable climate ceases to occur for 12 species in summer, potentially resulting in extirpation of those species from the Park (e.g., Figure 2). Climate is projected to become suitable in summer for 17 species not found at the Park today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 9, remain stable for 17, and worsen for 4 species. Suitable climate ceases to occur for 2 species in winter, potentially resulting in extirpation from the Park. Climate is projected to become suitable in winter for 37 species not found at the Park today, potentially resulting in local colonization.

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Park based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Park is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.




Figure 1. Projected changes in climate suitability for birds at the Park, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Park between the present and 2050 is 0.22 in summer (35th percentile across all national parks) and 0.19 in winter (25th percentile) under the high-emissions pathway. Potential species turnover declines to 0.12 in summer and 0.12 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Park is home to one species, the Brown-headed Nuthatch (*Sitta pusilla*), that is highly sensitive to climate change across its range (i.e., it is projected to lose climate suitability in over 50% of its current summer range in North America by 2050; Table 1; Langham et al. 2015).

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Abraham Lincoln Birthplace National Historical Park falls within the intermediate change group.** Parks anticipating intermediate change can best support landscape-scale bird conservation by emphasizing habitat restoration, maintaining natural

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

Suitable climate is not projected to disappear for this species at the Park; instead the Park may serve as an important refuge for this climate-sensitive species.



Figure 2. Although currently found at the Park, suitable climate for the American Goldfinch (*Spinus tristis*) may cease to occur here in summer by 2050, potentially resulting in local seasonal extirpation. Photo by John Benson/Flickr (CC BY 2.0).

disturbance regimes, and reducing other stressors. Furthermore, park managers have an opportunity to focus on supporting the 1 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Park based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Park is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend
Cackling/Canada Goose	-	Stable
Blue-winged Teal	-	Potential colonization
Bufflehead	-	Potential colonization
Northern Bobwhite	Stable	Stable
Eared Grebe	-	Potential colonization
Wood Stork	Potential colonization	-
Anhinga	-	Potential colonization
American White Pelican	-	Potential colonization
Great Egret	Potential colonization	Potential colonization
Little Blue Heron	Potential colonization	-
Cattle Egret	Potential colonization	-
Yellow-crowned Night-Heron	Potential colonization	-

Common Name	Summer Trend	Winter Trend
White Ibis	-	Potential colonization
Black Vulture	Improving	Improving
Mississippi Kite	Potential colonization	-
Northern Harrier	-	Stable
Red-tailed Hawk	Stable	Stable
Virginia Rail	-	Potential colonization
Spotted Sandpiper	-	Potential colonization
Greater Yellowlegs	-	Potential colonization
Least Sandpiper	-	Potential colonization
Gull-billed Tern	-	Potential colonization
Forster's Tern	-	Potential colonization
Eurasian Collared-Dove	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Mourning Dove	Stable	Worsening
Inca Dove	Potential colonization	Potential colonization
Yellow-billed Cuckoo	Improving	-
Greater Roadrunner	Potential colonization	-
Barn Owl	-	Potential colonization
Eastern Screech-Owl	x	Improving
Great Horned Owl	x	Potential extirpation
Barred Owl	x	Improving
Common Nighthawk	Improving	-
Chimney Swift	Stable	-
Red-bellied Woodpecker	Stable	Stable
Downy Woodpecker	Stable	Stable
Hairy Woodpecker	Potential extirpation	Worsening*
Red-cockaded Woodpecker	-	Potential colonization
Northern Flicker	Stable	Stable
Pileated Woodpecker	Stable	Stable
Eastern Wood-Pewee	Worsening	-
Acadian Flycatcher	Worsening	-
Eastern Phoebe	Stable	-
Scissor-tailed Flycatcher	Potential colonization	-
Loggerhead Shrike	Potential colonization	-
White-eyed Vireo	-	Potential colonization
Yellow-throated Vireo	Stable	-
Red-eyed Vireo	Potential extirpation	-
Blue Jay	Stable	Stable
American Crow	Stable	Stable
Barn Swallow	Worsening	-
Cliff Swallow	Potential colonization	-

Common Name	Summer Trend	Winter Trend
Carolina Chickadee	Improving	Improving
Tufted Titmouse	Stable	Improving
White-breasted Nuthatch	Potential extirpation	Worsening*
Brown-headed Nuthatch	Potential colonization^	-
House Wren	-	Potential colonization
Sedge Wren	-	Potential colonization
Marsh Wren	-	Potential colonization
Carolina Wren	Stable	Stable
Bewick's Wren	-	Potential colonization
Blue-gray Gnatcatcher	Stable	-
Golden-crowned Kinglet	-	Stable
Wood Thrush	Worsening*	-
American Robin	Potential extirpation	Stable
European Starling	Potential extirpation	Stable
American Pipit	-	Potential colonization
Sprague's Pipit	-	Potential colonization
Cedar Waxwing	Potential extirpation	Stable
Chestnut-collared Longspur	-	Potential colonization
Smith's Longspur	-	Potential colonization
Black-and-white Warbler	Improving	-
Swainson's Warbler	Potential colonization	-
Orange-crowned Warbler	-	Potential colonization
Common Yellowthroat	Potential extirpation	Potential colonization
Yellow-rumped Warbler	-	Improving
Prairie Warbler	Stable	-

Common Name	Summer Trend	Winter Trend
Yellow-breasted Chat	Stable	-
Eastern Towhee	Potential extirpation	x
Bachman's Sparrow	Potential colonization	Potential colonization
Chipping Sparrow	Potential extirpation	-
Vesper Sparrow	-	Potential colonization
Lark Sparrow	Potential colonization	-
Lincoln's Sparrow	-	Potential colonization
White-throated Sparrow	-	Stable
Harris's Sparrow	-	Potential colonization
Dark-eyed Junco	-	Worsening
Summer Tanager	Improving	-
Scarlet Tanager	Potential extirpation	-

Common Name	Summer Trend	Winter Trend
Northern Cardinal	Improving	Improving
Pyrrhuloxia	-	Potential colonization
Indigo Bunting	Improving	-
Painted Bunting	Potential colonization	-
Western Meadowlark	-	Potential colonization
Brewer's Blackbird	-	Potential colonization
Common Grackle	Worsening	Improving
Great-tailed Grackle	Potential colonization	Potential colonization
Bronzed Cowbird	-	Potential colonization
Brown-headed Cowbird	Worsening	Improving
House Finch	Potential extirpation	Potential extirpation
American Goldfinch	Potential extirpation	Stable