



Bent's Old Fort National Historic Site

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Bent's Old Fort National Historic Site (hereafter, the Site) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Site, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1). Among the species likely to be found at the Site today, climate suitability in summer under the high-emissions pathway is projected to improve for 10, remain stable for 24 (e.g., Figure 2), and worsen for 9 species. Suitable climate ceases to occur for 11 species in summer, potentially resulting in extirpation of those species from the Site. Climate is projected to become suitable in summer for 23 species not found at the Site today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 3, remain stable for 14, and worsen for 3 species. Suitable climate does not cease to occur for any species in winter. Climate is projected to become suitable in winter for 48 species not found at the Site today, potentially resulting in local colonization.

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Site based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Site is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.




Figure 1. Projected changes in climate suitability for birds at the Site, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Site between the present and 2050 is 0.26 in summer (44th percentile across all national parks) and 0.19 in winter (25th percentile) under the high-emissions pathway. Potential species turnover declines to 0.17 in summer and 0.08 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Site is or may become home to 7 species that are highly sensitive to climate change across their range (i.e., they are projected to lose climate suitability in over 50% of their current range in North America in summer and/or winter by 2050; Table 1; Langham et al. 2015). While the

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Bent's Old Fort National Historic Site falls within the high potential colonization group.** Parks anticipating high potential colonization can focus on actions that increase species' ability to respond to environmental change, such as increasing the amount of potential habitat, working with cooperating agencies and landowners to improve habitat connectivity for birds

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

Site may serve as an important refuge for 6 of these climate-sensitive species, one, the Northern Harrier (*Circus cyaneus*), might be extirpated from the Site in summer by 2050.



Figure 2. Climate at the Site in summer is projected to remain suitable for the Red-winged Blackbird (*Agelaius phoeniceus*) through 2050. Photo by Andy Reago & Chrissy McClarren/Flickr (CC BY 2.0).

across boundaries, managing the disturbance regime, and possibly more intensive management actions. Furthermore, park managers have an opportunity to focus on supporting the 6 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Site based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Site is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend
Mallard	Stable [^]	Stable
Blue-winged Teal	Stable	Potential colonization
Cinnamon Teal	x	Potential colonization
Northern Shoveler	Stable [^]	-
Scaled Quail	Stable	-
Gambel's Quail	Potential colonization	-
Northern Bobwhite	Improving*	Improving
Ring-necked Pheasant	Worsening	Improving
Wild Turkey	x	Stable
Least Grebe	-	Potential colonization
Horned Grebe	-	Potential colonization
Neotropic Cormorant	-	Potential colonization
American Bittern	-	Potential colonization [^]
Great Blue Heron	Potential extirpation	Stable

Common Name	Summer Trend	Winter Trend
Cattle Egret	Potential colonization	-
Yellow-crowned Night-Heron	Potential colonization	-
Mississippi Kite	Potential colonization	-
Northern Harrier	Potential extirpation [^]	-
Harris's Hawk	-	Potential colonization
Gray Hawk	Potential colonization	-
Swainson's Hawk	Stable [^]	-
Red-tailed Hawk	Stable	Improving
Common Gallinule	-	Potential colonization
American Coot	x	Stable
American Avocet	x	Potential colonization [^]
Killdeer	Worsening	Stable
Spotted Sandpiper	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Greater Yellowlegs	-	Potential colonization
Dunlin	-	Potential colonization [^]
Least Sandpiper	-	Potential colonization
Long-billed Dowitcher	-	Potential colonization
Bonaparte's Gull	-	Potential colonization
Gull-billed Tern	-	Potential colonization
Black Tern	Stable	-
Rock Pigeon	Stable	Stable
Mourning Dove	Stable	Stable
Yellow-billed Cuckoo	Improving	-
Greater Roadrunner	Potential colonization	-
Great Horned Owl	x	Worsening*
Burrowing Owl	-	Potential colonization
Lesser Nighthawk	Potential colonization	-
Common Nighthawk	Worsening	-
Chimney Swift	Stable	-
White-throated Swift	-	Potential colonization
Costa's Hummingbird	-	Potential colonization
Belted Kingfisher	Potential extirpation	Worsening*
Red-headed Woodpecker	Improving	-
Gila Woodpecker	Potential colonization	Potential colonization
Golden-fronted Woodpecker	-	Potential colonization
Red-naped Sapsucker	-	Potential colonization
Northern Flicker	Potential extirpation	Stable

Common Name	Summer Trend	Winter Trend
Gilded Flicker	Potential colonization	Potential colonization
American Kestrel	x	Stable
Gray Flycatcher	-	Potential colonization
Black Phoebe	-	Potential colonization
Vermilion Flycatcher	-	Potential colonization
Ash-throated Flycatcher	Potential colonization	-
Brown-crested Flycatcher	Potential colonization	-
Western Kingbird	Worsening	-
Eastern Kingbird	Potential extirpation	-
Scissor-tailed Flycatcher	Potential colonization	-
Loggerhead Shrike	Improving*	-
Warbling Vireo	Stable	-
Blue Jay	Stable	Stable
Horned Lark	Worsening*	-
Violet-green Swallow	Stable	Potential colonization
Barn Swallow	Worsening*	-
Cliff Swallow	Improving	-
Verdin	Potential colonization	Potential colonization
Rock Wren	Improving	-
House Wren	Potential extirpation	-
Cactus Wren	-	Potential colonization
Black-tailed Gnatcatcher	Potential colonization	-
Western Bluebird	-	Potential colonization
American Robin	Potential extirpation	Stable
Brown Thrasher	Stable	Potential colonization

Common Name	Summer Trend	Winter Trend
LeConte's Thrasher	-	Potential colonization
Crissal Thrasher	Potential colonization	Potential colonization
Northern Mockingbird	Stable	Stable
European Starling	Potential extirpation	-
Sprague's Pipit	-	Potential colonization
Phainopepla	Potential colonization	-
Smith's Longspur	-	Potential colonization
Lucy's Warbler	Potential colonization	-
Common Yellowthroat	Stable	-
Yellow Warbler	Potential extirpation	-
Yellow-breasted Chat	Stable	-
Abert's Towhee	Potential colonization	Potential colonization
Rufous-winged Sparrow	-	Potential colonization
Cassin's Sparrow	Stable	Potential colonization
Chipping Sparrow	-	Potential colonization
Field Sparrow	-	Potential colonization
Vesper Sparrow	-	Potential colonization
Lark Sparrow	Worsening*	-
Black-throated Sparrow	Potential colonization	Potential colonization
Lark Bunting	Worsening*	-
Savannah Sparrow	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Henslow's Sparrow	-	Potential colonization
Pyrrhuloxia	-	Potential colonization
Black-headed Grosbeak	Stable	-
Blue Grosbeak	Improving	-
Indigo Bunting	Improving	-
Painted Bunting	Potential colonization	-
Dickcissel	Stable	-
Red-winged Blackbird	Stable	Stable
Eastern Meadowlark	Potential colonization	Potential colonization
Western Meadowlark	Worsening	Worsening
Yellow-headed Blackbird	Stable	-
Common Grackle	Potential extirpation	-
Bronzed Cowbird	-	Potential colonization
Brown-headed Cowbird	Stable	-
Orchard Oriole	Potential extirpation	-
Hooded Oriole	Potential colonization	-
Bullock's Oriole	Improving	-
Scott's Oriole	Potential colonization	-
House Finch	Improving*	-
Lesser Goldfinch	-	Potential colonization
American Goldfinch	Stable	-
House Sparrow	x	Stable