



Blue Ridge Parkway

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Blue Ridge Parkway (hereafter, the Parkway) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Parkway, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1). Among the species likely to be found at the Parkway today, climate suitability in summer under the high-emissions pathway is projected to improve for 42 (e.g., Figure 2), remain stable for 18, and worsen for 15 species. Suitable climate ceases to occur for 44 species in summer, potentially resulting in extirpation of those species from the Parkway. Climate is projected to become suitable in summer for 19 species not found at the Parkway today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 55, remain stable for 11, and worsen for 12 species. Suitable climate ceases to occur for 7 species in winter, potentially resulting in extirpation from the Parkway. Climate is projected to become suitable in winter for 33 species not found at the Parkway today, potentially resulting in local colonization.

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Parkway based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Parkway is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.




Figure 1. Projected changes in climate suitability for birds at the Parkway, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Parkway between the present and 2050 is 0.23 in summer (38th percentile across all national parks) and 0.21 in winter (29th percentile) under the high-emissions pathway. Potential species turnover declines to 0.11 in summer and 0.10 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Parkway is or may become home to 12 species that are highly sensitive to climate change across their range (i.e., they are projected to lose climate suitability in over 50% of their current range in North America in summer and/or winter by 2050; Table 1; Langham et al. 2015). While the

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Blue Ridge Parkway falls within the high potential colonization group.** Parks anticipating high potential colonization can focus on actions that increase species' ability to respond to environmental change, such as increasing the amount of potential habitat, working with cooperating agencies and landowners to improve habitat

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

Parkway may serve as an important refuge for 9 of these climate-sensitive species, 3 might be extirpated from the Parkway in at least one season by 2050.



Figure 2. Climate at the Parkway in summer is projected to remain suitable for the Northern Cardinal (*Cardinalis cardinalis*) through 2050. Photo by Andy Morffew/Flickr (CC BY 2.0).

connectivity for birds across boundaries, managing the disturbance regime, and possibly more intensive management actions. Furthermore, park managers have an opportunity to focus on supporting the 9 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Parkway based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Parkway is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend
Cackling/Canada Goose	x	Worsening
Mute Swan	x	Potential extirpation
Wood Duck	x	Improving
Gadwall	-	Improving
Mallard	Potential extirpation [^]	Worsening
Blue-winged Teal	-	Potential colonization
Northern Shoveler	-	Potential colonization
Green-winged Teal	-	Potential colonization
Ring-necked Duck	x	Improving
Greater Scaup	-	Potential colonization [^]
Lesser Scaup	-	Potential colonization
Bufflehead	-	Improving
Hooded Merganser	x	Improving [^]

Common Name	Summer Trend	Winter Trend
Common Merganser	-	Potential extirpation
Red-breasted Merganser	-	Potential colonization [^]
Ruddy Duck	-	Potential colonization
Northern Bobwhite	Improving*	-
Ruffed Grouse	x	Potential extirpation
Wild Turkey	x	Worsening*
Common Loon	Potential extirpation	-
Pied-billed Grebe	x	Improving
Eared Grebe	-	Potential colonization
Double-crested Cormorant	x	Potential colonization
American White Pelican	-	Potential colonization
Great Blue Heron	Improving	Improving

Common Name	Summer Trend	Winter Trend
Great Egret	Improving	-
Little Blue Heron	Potential colonization	-
Cattle Egret	Potential colonization	-
Green Heron	Improving	-
Yellow-crowned Night-Heron	Potential colonization	-
Black Vulture	Improving	Improving
Turkey Vulture	x	Improving
Golden Eagle	x	Potential extirpation
Mississippi Kite	Potential colonization	-
Northern Harrier	Stable^	Improving
Sharp-shinned Hawk	x	Improving
Cooper's Hawk	x	Worsening*
Bald Eagle	x	Potential colonization
Red-shouldered Hawk	Improving*	Improving
Red-tailed Hawk	Improving	Improving
American Coot	-	Improving
Killdeer	Improving	Improving
Greater Yellowlegs	-	Potential colonization
Least Sandpiper	-	Potential colonization
American Woodcock	x	Improving
Pigeon Guillemot	Potential colonization	-
Bonaparte's Gull	-	Potential colonization
Laughing Gull	Potential colonization^	-
Ring-billed Gull	-	Improving
Forster's Tern	-	Potential colonization
Rock Pigeon	Stable	Worsening
Mourning Dove	Improving	Stable

Common Name	Summer Trend	Winter Trend
Yellow-billed Cuckoo	Improving*	-
Black-billed Cuckoo	Potential extirpation	-
Greater Roadrunner	Potential colonization	Potential colonization
Barn Owl	-	Potential colonization
Western Screech-Owl	-	Potential colonization
Eastern Screech-Owl	x	Improving
Great Horned Owl	x	Stable
Burrowing Owl	Potential colonization^	-
Barred Owl	x	Improving
Common Nighthawk	Improving	-
Chuck-will's-widow	Potential colonization	-
Chimney Swift	Stable	-
Ruby-throated Hummingbird	Improving	-
Belted Kingfisher	Stable	Improving
Red-headed Woodpecker	Improving*	Improving*
Red-bellied Woodpecker	Improving	Improving
Yellow-bellied Sapsucker	Potential extirpation	Improving
Downy Woodpecker	Improving	Stable
Hairy Woodpecker	Worsening	Worsening
Red-cockaded Woodpecker	-	Potential colonization
Northern Flicker	Potential extirpation	Improving
Pileated Woodpecker	Stable	Stable
American Kestrel	x	Improving
Eastern Wood-Pewee	Stable	-
Acadian Flycatcher	Worsening	-
Alder Flycatcher	Potential extirpation	-
Willow Flycatcher	Potential extirpation	-

Common Name	Summer Trend	Winter Trend
Least Flycatcher	Potential extirpation	-
Eastern Phoebe	Stable	Improving
Great Crested Flycatcher	Improving	-
Western Kingbird	Potential colonization	-
Eastern Kingbird	Improving	-
Scissor-tailed Flycatcher	Potential colonization	-
Loggerhead Shrike	Potential colonization	Potential colonization
White-eyed Vireo	Improving*	-
Bell's Vireo	Potential colonization	-
Yellow-throated Vireo	Stable	-
Warbling Vireo	Potential extirpation	-
Red-eyed Vireo	Worsening	-
Blue Jay	Improving	Worsening
American Crow	Worsening	Worsening
Fish Crow	Stable	Improving
Common Raven	Potential extirpation	Potential extirpation
Horned Lark	Stable	-
Northern Rough-winged Swallow	Improving	-
Purple Martin	Improving*	-
Tree Swallow	Potential extirpation	-
Barn Swallow	Improving	-
Cliff Swallow	Improving*	-
Carolina Chickadee	Improving	Improving
Black-capped Chickadee	Potential extirpation	Potential extirpation
Tufted Titmouse	Improving	Improving
Red-breasted Nuthatch	Potential extirpation	Stable
White-breasted Nuthatch	Worsening	Worsening
Brown-headed Nuthatch	Improving^	Improving*

Common Name	Summer Trend	Winter Trend
Brown Creeper	Potential extirpation^	Stable
House Wren	Potential extirpation	Potential colonization
Pacific/Winter Wren	-	Improving
Sedge Wren	-	Potential colonization
Carolina Wren	Improving	Improving
Blue-gray Gnatcatcher	Improving	-
Golden-crowned Kinglet	Potential extirpation	Stable
Ruby-crowned Kinglet	Potential extirpation	Improving
Eastern Bluebird	Improving	Improving
Veery	Potential extirpation	-
Swainson's Thrush	Potential extirpation	-
Hermit Thrush	Potential extirpation	Improving
Wood Thrush	Worsening*	-
American Robin	Potential extirpation	Improving
Gray Catbird	Potential extirpation	Improving
Brown Thrasher	Improving	Improving*
Northern Mockingbird	Improving	Improving
European Starling	Worsening	Worsening
American Pipit	-	Improving*
Cedar Waxwing	Potential extirpation	Improving
Smith's Longspur	-	Potential colonization
Ovenbird	Potential extirpation	-
Worm-eating Warbler	Worsening	-
Golden-winged Warbler	Stable	-
Black-and-white Warbler	Worsening	-
Prothonotary Warbler	Potential colonization	-

Common Name	Summer Trend	Winter Trend
Swainson's Warbler	Stable	-
Orange-crowned Warbler	-	Potential colonization
Mourning Warbler	Potential extirpation	-
Kentucky Warbler	Improving	-
Common Yellowthroat	Worsening	Potential colonization
Hooded Warbler	Stable	-
American Redstart	Potential extirpation	-
Northern Parula	Stable	-
Magnolia Warbler	Potential extirpation	-
Blackburnian Warbler	Potential extirpation	-
Yellow Warbler	Potential extirpation	-
Chestnut-sided Warbler	Potential extirpation	-
Black-throated Blue Warbler	Potential extirpation	-
Pine Warbler	Improving*^	Improving*
Yellow-rumped Warbler	Potential extirpation	Improving
Yellow-throated Warbler	Stable	-
Prairie Warbler	Improving	-
Black-throated Green Warbler	Potential extirpation	-
Canada Warbler	Potential extirpation	-
Yellow-breasted Chat	Improving	-
Eastern Towhee	Stable	x
Bachman's Sparrow	Potential colonization	Potential colonization
American Tree Sparrow	-	Potential extirpation
Chipping Sparrow	Worsening	Improving*
Field Sparrow	Stable	Improving

Common Name	Summer Trend	Winter Trend
Vesper Sparrow	Potential extirpation	Potential colonization
Lark Sparrow	Potential colonization	-
Savannah Sparrow	Potential extirpation	Potential colonization
Grasshopper Sparrow	Improving	-
Henslow's Sparrow	-	Potential colonization
LeConte's Sparrow	-	Potential colonization
Seaside Sparrow	Potential colonization^	-
Fox Sparrow	-	Improving
Song Sparrow	Potential extirpation	Improving
Lincoln's Sparrow	-	Potential colonization
Swamp Sparrow	-	Improving
White-throated Sparrow	Potential extirpation	Improving
Harris's Sparrow	-	Potential colonization
White-crowned Sparrow	-	Stable
Dark-eyed Junco	x	Stable
Summer Tanager	Improving*	-
Scarlet Tanager	Worsening*	-
Northern Cardinal	Improving	Improving
Rose-breasted Grosbeak	Potential extirpation	-
Blue Grosbeak	Improving*	-
Indigo Bunting	Improving	-
Painted Bunting	Potential colonization	-
Dickcissel	Potential colonization	-
Bobolink	Potential extirpation	-
Red-winged Blackbird	Worsening	Improving
Eastern Meadowlark	Improving*	Improving*

Common Name	Summer Trend	Winter Trend
Brewer's Blackbird	-	Potential colonization
Common Grackle	Worsening	Improving
Brown-headed Cowbird	Stable	Improving
Orchard Oriole	Improving*	-
Baltimore Oriole	Potential extirpation	-
House Finch	Potential extirpation	Worsening*

Common Name	Summer Trend	Winter Trend
Purple Finch	-	Stable
Red Crossbill	Potential extirpation [^]	-
Pine Siskin	Potential extirpation	Improving
American Goldfinch	Worsening*	Stable
House Sparrow	x	Worsening*