



Carl Sandburg Home National Historic Site

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Carl Sandburg Home National Historic Site (hereafter, the Site) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Site, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1). Among the species likely to be found at the Site today, climate suitability in summer under the high-emissions pathway is projected to improve for 23, remain stable for 16, and worsen for 14 species. Suitable climate ceases to occur for 12 species in summer, potentially resulting in extirpation of those species from the Site (e.g., Figure 2). Climate is projected to become suitable in summer for 13 species not found at the Site today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 29, remain stable for 15, and worsen for 4 species. Suitable climate does not cease to occur for any species in winter. Climate is projected to become suitable in winter for 34 species not found at the Site today, potentially resulting in local colonization.

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Site based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Site is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.




Figure 1. Projected changes in climate suitability for birds at the Site, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Site between the present and 2050 is 0.16 in summer (23rd percentile across all national parks) and 0.18 in winter (24th percentile) under the high-emissions pathway. Potential species turnover declines to 0.09 in summer and 0.08 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Site is or may become home to 6 species that are highly sensitive to climate change across their range (i.e., they are projected to lose climate suitability in over 50% of their current range in North America in summer and/or winter by 2050; Table 1; Langham et al. 2015). Suitable climate is

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Carl Sandburg Home National Historic Site falls within the low change group.** Parks anticipating low change can best support landscape-scale bird conservation by emphasizing habitat restoration, maintaining natural disturbance regimes, and reducing

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

not projected to disappear for these 6 species at the Site; instead the Site may serve as an important refuge for these climate-sensitive species.



Figure 2. Although currently found at the Site, suitable climate for the American Goldfinch (*Spinus tristis*) may cease to occur here in summer by 2050, potentially resulting in local seasonal extirpation. Photo by John Benson/Flickr (CC BY 2.0).

other stressors. Furthermore, park managers have an opportunity to focus on supporting the 6 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Site based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Site is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend
Cackling/Canada Goose	x	Worsening*
Gadwall	-	Improving
American Wigeon	-	Stable
Mallard	Stable^	Stable
Blue-winged Teal	-	Potential colonization
Northern Shoveler	-	Potential colonization
Hooded Merganser	-	Improving^
Red-breasted Merganser	-	Potential colonization^
Pied-billed Grebe	-	Improving
Eared Grebe	-	Potential colonization
Wood Stork	Potential colonization	-
Double-crested Cormorant	-	Potential colonization
Great Blue Heron	Improving	Improving
Great Egret	Potential colonization	Potential colonization

Common Name	Summer Trend	Winter Trend
Little Blue Heron	Potential colonization	-
Cattle Egret	-	Potential colonization
Green Heron	Improving	-
Black Vulture	Potential colonization	-
Turkey Vulture	x	Improving
Osprey	-	Potential colonization
Mississippi Kite	Potential colonization	-
Northern Harrier	-	Potential colonization
Sharp-shinned Hawk	-	Worsening*
Cooper's Hawk	x	Stable
Red-tailed Hawk	Improving	Improving
Sora	-	Potential colonization
Spotted Sandpiper	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Least Sandpiper	-	Potential colonization
Pigeon Guillemot	Potential colonization	-
Bonaparte's Gull	-	Potential colonization
Ring-billed Gull	-	Potential colonization
Herring Gull	-	Potential colonization [^]
Gull-billed Tern	-	Potential colonization
Forster's Tern	-	Potential colonization
Eurasian Collared-Dove	-	Potential colonization
Mourning Dove	Stable	Improving
Yellow-billed Cuckoo	Improving	-
Greater Roadrunner	-	Potential colonization
Barn Owl	-	Potential colonization
Chimney Swift	Stable	-
Allen's Hummingbird	-	Potential colonization
Belted Kingfisher	Improving	Improving
Red-headed Woodpecker	Improving*	Improving*
Red-bellied Woodpecker	Improving	Stable
Yellow-bellied Sapsucker	-	Improving
Downy Woodpecker	Stable	Improving
Hairy Woodpecker	Stable	Stable
Red-cockaded Woodpecker	-	Potential colonization
Northern Flicker	Stable	Stable
Pileated Woodpecker	Worsening	Improving
American Kestrel	x	Improving
Eastern Wood-Pewee	Worsening	-
Acadian Flycatcher	Stable	-
Eastern Phoebe	Worsening	-

Common Name	Summer Trend	Winter Trend
Great Crested Flycatcher	Improving*	-
Eastern Kingbird	Improving	-
Loggerhead Shrike	Potential colonization	-
White-eyed Vireo	Improving	Potential colonization
Yellow-throated Vireo	Worsening	-
Red-eyed Vireo	Worsening	-
Blue Jay	Improving	Stable
American Crow	Stable	Stable
Northern Rough-winged Swallow	Improving	-
Purple Martin	Improving*	-
Tree Swallow	Potential extirpation	-
Barn Swallow	Improving	-
Cliff Swallow	Potential colonization	-
Carolina Chickadee	Stable	Improving
Tufted Titmouse	Worsening	Improving
White-breasted Nuthatch	Worsening*	Worsening*
Brown-headed Nuthatch	Potential colonization [^]	-
Brown Creeper	-	Stable
House Wren	Potential extirpation	-
Sedge Wren	-	Potential colonization
Carolina Wren	Improving	Improving
Blue-gray Gnatcatcher	Stable	Potential colonization
Golden-crowned Kinglet	-	Improving
Ruby-crowned Kinglet	-	Improving
Eastern Bluebird	Improving	Improving
Hermit Thrush	-	Improving
Wood Thrush	Stable	-
American Robin	Potential extirpation	Improving

Common Name	Summer Trend	Winter Trend
Gray Catbird	Potential extirpation	Potential colonization
Brown Thrasher	Stable	-
Northern Mockingbird	Improving	Improving
Cedar Waxwing	Potential extirpation	Improving
Ovenbird	Potential extirpation	-
Worm-eating Warbler	Worsening	-
Blue-winged Warbler	Worsening	-
Black-and-white Warbler	Worsening	Potential colonization
Prothonotary Warbler	Potential colonization	-
Orange-crowned Warbler	-	Potential colonization
Common Yellowthroat	-	Potential colonization
Hooded Warbler	Stable	-
American Redstart	Potential extirpation	-
Yellow Warbler	Potential extirpation	-
Pine Warbler	Improving^	Improving*
Yellow-rumped Warbler	-	Stable
Yellow-throated Warbler	Worsening	-
Black-throated Green Warbler	Potential extirpation	-
Yellow-breasted Chat	Improving	-
Eastern Towhee	Stable	x
Bachman's Sparrow	Potential colonization	Potential colonization

Common Name	Summer Trend	Winter Trend
Chipping Sparrow	Worsening	-
Vesper Sparrow	-	Potential colonization
LeConte's Sparrow	-	Potential colonization
Fox Sparrow	-	Improving
Song Sparrow	Potential extirpation	Stable
Lincoln's Sparrow	-	Potential colonization
Swamp Sparrow	-	Improving
White-throated Sparrow	-	Stable
Dark-eyed Junco	x	Worsening
Scarlet Tanager	Worsening*	-
Northern Cardinal	Improving	Stable
Blue Grosbeak	Improving*	-
Indigo Bunting	Improving	-
Painted Bunting	Potential colonization	-
Red-winged Blackbird	Potential extirpation	Improving
Brewer's Blackbird	-	Potential colonization
Common Grackle	Worsening	Improving
Great-tailed Grackle	Potential colonization	-
Brown-headed Cowbird	Stable	Improving*
Orchard Oriole	Improving	-
House Finch	Stable	Stable
American Goldfinch	Potential extirpation	Stable