



Catoctin Mountain Park

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Catoctin Mountain Park (hereafter, the Park) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Park, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1). Among the species likely to be found at the Park today, climate suitability in summer under the high-emissions pathway is projected to improve for 35 (e.g., Figure 2), remain stable for 13, and worsen for 10 species. Suitable climate ceases to occur for 18 species in summer, potentially resulting in extirpation of those species from the Park. Climate is projected to become suitable in summer for 19 species not found at the Park today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 25, remain stable for 5, and worsen for 12 species. Suitable climate ceases to occur for 2 species in winter, potentially resulting in extirpation from the Park. Climate is projected to become suitable in winter for 34 species not found at the Park today, potentially resulting in local colonization.

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Park based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Park is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.




Figure 1. Projected changes in climate suitability for birds at the Park, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Park between the present and 2050 is 0.23 in summer (37th percentile across all national parks) and 0.19 in winter (24th percentile) under the high-emissions pathway. Potential species turnover declines to 0.14 in summer and 0.12 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Park is or may become home to 4 species that are highly sensitive to climate change across their range (i.e., they are projected to lose climate suitability in over 50% of their current range in North America in summer and/or winter by 2050; Table 1; Langham et al. 2015). While the

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Catoctin Mountain Park falls within the intermediate change group.** Parks anticipating intermediate change can best support landscape-scale bird conservation by emphasizing habitat restoration, maintaining natural disturbance regimes, and reducing

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

Park may serve as an important refuge for 3 of these climate-sensitive species, one, the Mallard (*Anas platyrhynchos*), might be extirpated from the Park in summer by 2050.



Figure 2. Climate at the Park in summer is projected to remain suitable for the Northern Cardinal (*Cardinalis cardinalis*) through 2050. Photo by Andy Morffew/Flickr (CC BY 2.0).

other stressors. Furthermore, park managers have an opportunity to focus on supporting the 3 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Park based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Park is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend
Cackling/Canada Goose	x	Worsening
Mallard	Potential extirpation [^]	Stable
Blue-winged Teal	-	Potential colonization
Green-winged Teal	-	Potential colonization
Bufflehead	-	Potential colonization
Red-breasted Merganser	-	Potential colonization [^]
Northern Bobwhite	Improving*	Improving
Eared Grebe	-	Potential colonization
American White Pelican	-	Potential colonization
Great Blue Heron	Improving	-
Great Egret	Potential colonization	-
Little Blue Heron	Potential colonization	-

Common Name	Summer Trend	Winter Trend
Cattle Egret	Potential colonization	-
Yellow-crowned Night-Heron	Potential colonization	-
Black Vulture	Improving	Improving
Turkey Vulture	x	Improving
Mississippi Kite	Potential colonization	-
Northern Harrier	-	Improving
Sharp-shinned Hawk	-	Improving
Bald Eagle	-	Potential colonization
Red-shouldered Hawk	Improving	Improving
Red-tailed Hawk	Improving	Improving
Killdeer	Improving	-
Least Sandpiper	-	Potential colonization
Bonaparte's Gull	-	Potential colonization
Forster's Tern	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Rock Pigeon	Worsening	Worsening*
Eurasian Collared-Dove	-	Potential colonization
Mourning Dove	Improving	Worsening
Yellow-billed Cuckoo	Improving*	-
Greater Roadrunner	Potential colonization	Potential colonization
Barn Owl	-	Potential colonization
Western Screech-Owl	-	Potential colonization
Common Nighthawk	Potential colonization	-
Chuck-will's-widow	Potential colonization	-
Chimney Swift	Worsening	-
Belted Kingfisher	Stable	-
Red-headed Woodpecker	Improving	Improving
Red-bellied Woodpecker	Improving	Improving
Yellow-bellied Sapsucker	-	Improving
Downy Woodpecker	Improving	Worsening
Hairy Woodpecker	Stable	Worsening*
Northern Flicker	Potential extirpation	Improving
Pileated Woodpecker	Improving	Improving
American Kestrel	-	Improving
Eastern Wood-Pewee	Improving	-
Acadian Flycatcher	Stable	-
Eastern Phoebe	Improving	Potential colonization
Great Crested Flycatcher	Improving	-
Eastern Kingbird	Stable	-
Scissor-tailed Flycatcher	Potential colonization	-
Loggerhead Shrike	Potential colonization	Potential colonization
White-eyed Vireo	Improving*	-
Bell's Vireo	Potential colonization	-

Common Name	Summer Trend	Winter Trend
Yellow-throated Vireo	Stable	-
Warbling Vireo	Stable	-
Red-eyed Vireo	Potential extirpation	-
Blue Jay	Improving	Stable
American Crow	Worsening	Worsening
Common Raven	Potential extirpation	Potential extirpation
Northern Rough-winged Swallow	Improving	-
Purple Martin	Improving*	-
Tree Swallow	Potential extirpation	-
Barn Swallow	Improving	-
Cliff Swallow	Potential colonization	-
Carolina Chickadee	Improving*	Improving
Tufted Titmouse	Improving	Improving
White-breasted Nuthatch	Stable	Worsening*
Brown-headed Nuthatch	Potential colonization^	Potential colonization
Brown Creeper	-	Stable
House Wren	Potential extirpation	Potential colonization
Sedge Wren	-	Potential colonization
Carolina Wren	Improving	Improving
Blue-gray Gnatcatcher	Improving*	-
Golden-crowned Kinglet	-	Improving
Eastern Bluebird	Improving	Improving
Veery	Potential extirpation	-
Wood Thrush	Worsening*	-
American Robin	Worsening	Improving
Gray Catbird	Potential extirpation	-
Brown Thrasher	Stable	Potential colonization
Northern Mockingbird	Improving	Improving

Common Name	Summer Trend	Winter Trend
European Starling	Worsening	Worsening
American Pipit	-	Potential colonization
Cedar Waxwing	Potential extirpation	-
Chestnut-collared Longspur	-	Potential colonization
Smith's Longspur	-	Potential colonization
Ovenbird	Potential extirpation	-
Worm-eating Warbler	Stable	-
Black-and-white Warbler	Improving	-
Prothonotary Warbler	Potential colonization	-
Swainson's Warbler	Potential colonization	-
Orange-crowned Warbler	-	Potential colonization
Kentucky Warbler	Improving	-
Common Yellowthroat	Worsening	-
Hooded Warbler	Stable	-
American Redstart	Potential extirpation	-
Northern Parula	Improving*	-
Yellow Warbler	Potential extirpation	-
Palm Warbler	-	Potential colonization ^
Pine Warbler	-	Potential colonization
Eastern Towhee	Stable	-
American Tree Sparrow	-	Worsening*
Chipping Sparrow	Potential extirpation	Improving
Field Sparrow	Improving	-
Vesper Sparrow	Potential extirpation	Potential colonization
Lark Sparrow	Potential colonization	-

Common Name	Summer Trend	Winter Trend
Savannah Sparrow	-	Potential colonization
Henslow's Sparrow	-	Potential colonization
LeConte's Sparrow	-	Potential colonization
Song Sparrow	Potential extirpation	Stable
Lincoln's Sparrow	-	Potential colonization
White-throated Sparrow	-	Improving
Harris's Sparrow	-	Potential colonization
Dark-eyed Junco	-	Stable
Summer Tanager	Improving*	-
Scarlet Tanager	Potential extirpation	-
Northern Cardinal	Improving	Improving
Rose-breasted Grosbeak	Potential extirpation	-
Blue Grosbeak	Potential colonization	-
Indigo Bunting	Improving	-
Painted Bunting	Potential colonization	-
Red-winged Blackbird	Stable	-
Eastern Meadowlark	Improving	-
Brewer's Blackbird	-	Potential colonization
Common Grackle	Worsening	Improving
Great-tailed Grackle	Potential colonization	Potential colonization
Brown-headed Cowbird	Stable	Improving
Orchard Oriole	Improving	-
Baltimore Oriole	Worsening*	-
House Finch	Potential extirpation	Worsening*
American Goldfinch	Worsening	Worsening
Evening Grosbeak	-	Potential extirpation

Common Name	Summer Trend	Winter Trend
House Sparrow	x	Worsening*