



Chattahoochee River National Recreation Area

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Chattahoochee River National Recreation Area (hereafter, the Recreation Area) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Recreation Area, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1).

Among the species likely to be found at the Recreation Area today, climate suitability in summer under the high-emissions pathway is projected to improve for 18, remain stable for 34, and worsen for 17 species. Suitable climate ceases to occur for 23 species in summer, potentially resulting in extirpation of those species from the Recreation Area (e.g., Figure 2). Climate is projected to become suitable in summer for 20 species not found at the Recreation Area today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 39, remain stable for 56, and worsen for 9 species. Suitable climate ceases to occur for 12 species in winter, potentially resulting in extirpation from the Recreation Area. Climate is projected to become suitable in winter for 40 species not

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Recreation Area based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Recreation Area is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.

found at the Recreation Area today, potentially resulting in local colonization.




Figure 1. Projected changes in climate suitability for birds at the Recreation Area, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Recreation Area between the present and 2050 is 0.23 in summer (38th percentile across all national parks) and 0.23 in winter (34th percentile) under the high-emissions pathway. Potential species turnover declines to 0.14 in summer and 0.14 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Recreation Area is or may become home to 13 species that are highly sensitive to climate change across their range (i.e., they are projected to lose climate suitability in over 50% of their current range in North America in summer and/or winter by 2050; Table 1; Langham et al. 2015). While the Recreation Area may serve as an

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Chattahoochee River National Recreation Area falls within the intermediate change group.** Parks anticipating intermediate change can best support landscape-scale bird conservation by emphasizing habitat restoration, maintaining natural

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

important refuge for 9 of these climate-sensitive species, 4 might be extirpated from the Recreation Area in at least one season by 2050.



Figure 2. Although currently found at the Recreation Area, suitable climate for the American Goldfinch (*Spinus tristis*) may cease to occur here in summer by 2050, potentially resulting in local seasonal extirpation. Photo by John Benson/Flickr (CC BY 2.0).

disturbance regimes, and reducing other stressors.

Furthermore, park managers have an opportunity to focus on supporting the 9 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Recreation Area based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Recreation Area is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend
Black-bellied Whistling-Duck	Potential colonization	-
Cackling/Canada Goose	x	Potential extirpation
Wood Duck	x	Stable
Gadwall	-	Improving
American Wigeon	-	Improving
American Black Duck	-	Potential extirpation
Mallard	Potential extirpation [^]	Stable
Mottled Duck	-	Potential colonization
Blue-winged Teal	-	Potential colonization
Northern Shoveler	-	Improving
Green-winged Teal	-	Stable
Ring-necked Duck	-	Improving
Greater Scaup	-	Stable [^]
Lesser Scaup	-	Improving

Common Name	Summer Trend	Winter Trend
White-winged Scoter	-	Potential extirpation
Bufflehead	-	Potential extirpation
Hooded Merganser	x	Potential extirpation [^]
Red-breasted Merganser	-	Potential extirpation [^]
Ruddy Duck	-	Stable
Northern Bobwhite	Worsening	-
Wild Turkey	x	Potential extirpation
Common Loon	-	Stable [^]
Pied-billed Grebe	x	Improving
Horned Grebe	-	Stable
Red-necked Grebe	-	Stable [^]
Eared Grebe	-	Improving
Wood Stork	Potential colonization	-

Common Name	Summer Trend	Winter Trend
Neotropic Cormorant	-	Potential colonization
Double-crested Cormorant	x	Improving
Anhinga	Potential colonization [^]	Potential colonization
American White Pelican	-	Potential colonization
Great Blue Heron	Stable	Stable
Great Egret	Improving*	Improving*
Snowy Egret	x	Potential colonization
Little Blue Heron	Improving*	Potential colonization
Cattle Egret	Improving*	Potential colonization
Green Heron	Stable	-
Yellow-crowned Night-Heron	Improving	-
White Ibis	Improving	Potential colonization
White-faced Ibis	-	Potential colonization [^]
Black Vulture	Improving	Stable
Turkey Vulture	x	Stable
Osprey	x	Improving
Mississippi Kite	Improving	-
Northern Harrier	-	Stable
Sharp-shinned Hawk	x	Stable
Cooper's Hawk	x	Worsening
Bald Eagle	x	Stable
Red-shouldered Hawk	Stable	Stable
Red-tailed Hawk	Stable	Improving
Ferruginous Hawk	-	Potential colonization
Virginia Rail	-	Stable
Sora	-	Potential colonization
American Coot	-	Improving

Common Name	Summer Trend	Winter Trend
Black-necked Stilt	-	Potential colonization
Killdeer	Potential extirpation	Improving
Spotted Sandpiper	x	Improving*
Greater Yellowlegs	-	Potential colonization
Lesser Yellowlegs	-	Potential colonization
Long-billed Curlew	-	Potential colonization
Stilt Sandpiper	-	Potential colonization
Least Sandpiper	-	Improving
Long-billed Dowitcher	-	Potential colonization
Wilson's Snipe	-	Stable
American Woodcock	x	Stable
Bonaparte's Gull	-	Stable
Ring-billed Gull	-	Worsening*
Herring Gull	-	Potential extirpation [^]
Rock Pigeon	Potential extirpation	Stable
Eurasian Collared-Dove	-	Potential colonization
White-winged Dove	-	Potential colonization
Mourning Dove	Stable	Worsening
Inca Dove	Potential colonization	Potential colonization
Common Ground-Dove	Potential colonization	Potential colonization
Yellow-billed Cuckoo	Improving*	-
Greater Roadrunner	Potential colonization	-
Eastern Screech-Owl	x	Improving
Great Horned Owl	x	Potential extirpation
Barred Owl	x	Improving

Common Name	Summer Trend	Winter Trend
Common Nighthawk	Potential colonization	-
Common Pauraque	-	Potential colonization
Chimney Swift	Worsening	-
Ruby-throated Hummingbird	Stable	-
Belted Kingfisher	Stable	Stable
Red-headed Woodpecker	Stable	Stable
Red-bellied Woodpecker	Stable	Stable
Yellow-bellied Sapsucker	-	Improving
Ladder-backed Woodpecker	Potential colonization	-
Downy Woodpecker	Worsening	Stable
Hairy Woodpecker	Potential extirpation	Worsening
Northern Flicker	Improving	Stable
Pileated Woodpecker	Improving*	Stable
Crested Caracara	Potential colonization	Potential colonization
American Kestrel	x	Worsening
Merlin	-	Improving^
Eastern Wood-Pewee	Worsening	-
Acadian Flycatcher	Stable	-
Eastern Phoebe	Worsening	Stable
Vermilion Flycatcher	Potential colonization	Potential colonization
Great Crested Flycatcher	Worsening	-
Brown-crested Flycatcher	Potential colonization	-
Eastern Kingbird	Worsening	-
Scissor-tailed Flycatcher	Potential colonization	-
White-eyed Vireo	Improving	Potential colonization
Yellow-throated Vireo	Stable	-
Red-eyed Vireo	Stable	-
Blue Jay	Stable	Stable

Common Name	Summer Trend	Winter Trend
American Crow	Stable	Stable
Fish Crow	Stable	Improving*
Northern Rough-winged Swallow	Stable	Potential colonization
Purple Martin	Improving	-
Tree Swallow	Potential extirpation	-
Barn Swallow	Worsening	-
Cliff Swallow	Improving*	-
Cave Swallow	Potential colonization	-
Carolina Chickadee	Improving	Improving
Tufted Titmouse	Worsening	Stable
Black-crested Titmouse	Potential colonization	-
Red-breasted Nuthatch	-	Stable
White-breasted Nuthatch	Potential extirpation	Potential extirpation
Brown-headed Nuthatch	Stable^	Stable
Brown Creeper	-	Stable
House Wren	Potential extirpation	Stable
Pacific/Winter Wren	-	Stable
Sedge Wren	-	Improving*
Marsh Wren	-	Potential colonization
Carolina Wren	Stable	Improving
Bewick's Wren	-	Potential colonization
Blue-gray Gnatcatcher	Worsening	Improving
Golden-crowned Kinglet	-	Stable
Ruby-crowned Kinglet	-	Improving
Eastern Bluebird	Worsening	Improving
Hermit Thrush	-	Stable
Wood Thrush	Worsening	-
American Robin	Potential extirpation	Stable

Common Name	Summer Trend	Winter Trend
Gray Catbird	Potential extirpation	Improving
Brown Thrasher	Worsening*	Stable
Northern Mockingbird	Worsening	Improving
European Starling	Potential extirpation	Worsening
American Pipit	-	Improving
Sprague's Pipit	-	Potential colonization
Cedar Waxwing	Potential extirpation	Stable
Chestnut-collared Longspur	-	Potential colonization
Smith's Longspur	-	Potential colonization
Ovenbird	Potential extirpation	-
Worm-eating Warbler	Stable	-
Blue-winged Warbler	Stable	-
Black-and-white Warbler	Stable	Stable
Prothonotary Warbler	Stable	-
Orange-crowned Warbler	-	Improving*
Kentucky Warbler	Improving*	-
Common Yellowthroat	Potential extirpation	Stable
Hooded Warbler	Improving*	-
American Redstart	Improving	-
Northern Parula	Stable	-
Yellow Warbler	Potential extirpation	-
Black-throated Blue Warbler	Stable	-
Palm Warbler	-	Worsening**^
Pine Warbler	Stable^	Stable
Yellow-rumped Warbler	-	Stable
Yellow-throated Warbler	Stable	-
Prairie Warbler	Stable	-
Yellow-breasted Chat	Stable	-

Common Name	Summer Trend	Winter Trend
Eastern Towhee	Worsening*	x
Cassin's Sparrow	-	Potential colonization
Bachman's Sparrow	Potential colonization	Potential colonization
Chipping Sparrow	Potential extirpation	Improving
Field Sparrow	Worsening*	Stable
Vesper Sparrow	-	Improving*
Lark Sparrow	Potential colonization	Potential colonization
Savannah Sparrow	Stable	Stable
Grasshopper Sparrow	Potential extirpation	Improving
Henslow's Sparrow	-	Potential colonization
LeConte's Sparrow	-	Improving*
Fox Sparrow	-	Worsening*
Song Sparrow	Potential extirpation	Stable
Lincoln's Sparrow	-	Improving*
Swamp Sparrow	-	Stable
White-throated Sparrow	-	Improving
Harris's Sparrow	-	Improving*
White-crowned Sparrow	-	Stable
Dark-eyed Junco	-	Worsening
Summer Tanager	Stable	-
Scarlet Tanager	Potential extirpation	-
Northern Cardinal	Improving	Stable
Pyrrhuloxia	-	Potential colonization
Blue Grosbeak	Worsening*	-
Indigo Bunting	Stable	-
Painted Bunting	Potential colonization	-
Red-winged Blackbird	Potential extirpation	Stable
Eastern Meadowlark	Stable	Stable

Common Name	Summer Trend	Winter Trend
Western Meadowlark	-	Potential colonization
Rusty Blackbird	-	Stable
Common Grackle	Potential extirpation	Improving
Great-tailed Grackle	Potential colonization	Potential colonization
Bronzed Cowbird	Potential colonization	Potential colonization
Brown-headed Cowbird	Potential extirpation	Improving
Orchard Oriole	Stable	-

Common Name	Summer Trend	Winter Trend
Hooded Oriole	Potential colonization	-
Baltimore Oriole	Stable	x
House Finch	Potential extirpation	Potential extirpation
Purple Finch	-	Potential extirpation
Pine Siskin	-	Stable
American Goldfinch	Potential extirpation	Stable
House Sparrow	x	Stable