



Fort Necessity National Battlefield

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Fort Necessity National Battlefield (hereafter, the Battlefield) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Battlefield, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1). Among the species likely to be found at the Battlefield today, climate suitability in summer under the high-emissions pathway is projected to improve for 18 (e.g., Figure 2), remain stable for 7, and worsen for 16 species. Suitable climate ceases to occur for 13 species in summer, potentially resulting in extirpation of those species from the Battlefield. Climate is projected to become suitable in summer for 21 species not found at the Battlefield today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 17, remain stable for 6, and worsen for 10 species. Suitable climate ceases to occur for 2 species in winter, potentially resulting in extirpation from the Battlefield. Climate is projected to become suitable in winter for 26 species not found at the Battlefield today, potentially resulting in local colonization.

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Battlefield based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Battlefield is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.




Figure 1. Projected changes in climate suitability for birds at the Battlefield, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Battlefield between the present and 2050 is 0.27 in summer (45th percentile across all national parks) and 0.24 in winter (34th percentile) under the high-emissions pathway. Potential species turnover declines to 0.20 in summer and 0.19 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Battlefield is or may become home to 3 species that are highly sensitive to climate change across their range (i.e., they are projected to lose climate suitability in over 50% of their current range in North America in summer and/or winter by 2050; Table 1; Langham et al. 2015).

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Fort Necessity National Battlefield falls within the high potential extirpation group.** Parks anticipating high potential extirpation can focus on actions that increase species' ability to respond to environmental change, such as increasing the amount of potential habitat, working with cooperating agencies and landowners to

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

While the Battlefield may serve as an important refuge for 2 of these climate-sensitive species, one, the Brown Creeper (*Certhia americana*), might be extirpated from the Battlefield in summer by 2050.



Figure 2. Climate at the Battlefield in summer is projected to remain suitable for the Northern Cardinal (*Cardinalis cardinalis*) through 2050. Photo by Andy Morffew/Flickr (CC BY 2.0).

improve habitat connectivity for birds across boundaries, managing the disturbance regime, and possibly more intensive management actions. Furthermore, park managers have an opportunity to focus on supporting the 2 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Battlefield based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Battlefield is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend
Cackling/Canada Goose	x	Stable
Wood Duck	x	Potential colonization
Mallard	-	Worsening
Northern Shoveler	-	Potential colonization
Green-winged Teal	-	Potential colonization
Lesser Scaup	-	Potential colonization
Bufflehead	-	Potential colonization
Red-breasted Merganser	-	Potential colonization [^]
Northern Bobwhite	Potential colonization	Potential colonization
Ring-necked Pheasant	-	Potential extirpation
Wild Turkey	x	Worsening*
Double-crested Cormorant	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Great Blue Heron	Stable	-
Great Egret	Potential colonization	-
Black Vulture	Potential colonization	Potential colonization
Turkey Vulture	x	Improving*
Mississippi Kite	Potential colonization	-
Cooper's Hawk	-	Stable
Bald Eagle	-	Improving
Red-shouldered Hawk	Potential colonization	-
Red-tailed Hawk	Improving	Improving
American Coot	-	Potential colonization
Killdeer	-	Improving
American Woodcock	-	Potential colonization
Mourning Dove	Improving	Worsening
Eastern Screech-Owl	-	Improving

Common Name	Summer Trend	Winter Trend
Great Horned Owl	-	Potential colonization
Chuck-will's-widow	Potential colonization	-
Chimney Swift	Stable	-
Red-bellied Woodpecker	Improving	Improving
Downy Woodpecker	Improving	Worsening
Hairy Woodpecker	Worsening	Worsening*
Northern Flicker	Potential extirpation	-
Pileated Woodpecker	Improving	Stable
Eastern Wood-Pewee	Improving	-
Acadian Flycatcher	Improving	-
Eastern Phoebe	Stable	Potential colonization
Eastern Kingbird	Improving	-
Scissor-tailed Flycatcher	Potential colonization	-
Loggerhead Shrike	-	Potential colonization
White-eyed Vireo	Potential colonization	-
Bell's Vireo	Potential colonization	-
Red-eyed Vireo	Worsening	-
Blue Jay	Stable	Stable
American Crow	Worsening	Worsening
Fish Crow	Potential colonization	-
Common Raven	Potential extirpation	-
Purple Martin	Potential colonization	-
Tree Swallow	Potential extirpation	-
Barn Swallow	Improving	-
Cliff Swallow	Potential colonization	-
Carolina Chickadee	Potential colonization	-

Common Name	Summer Trend	Winter Trend
Black-capped Chickadee	Potential extirpation	Potential extirpation
Tufted Titmouse	Improving	Improving
White-breasted Nuthatch	Stable	Worsening
Brown-headed Nuthatch	-	Potential colonization
Brown Creeper	Potential extirpation [^]	Improving
House Wren	Potential extirpation	-
Carolina Wren	-	Improving
Blue-gray Gnatcatcher	Improving	-
Golden-crowned Kinglet	-	Improving
Ruby-crowned Kinglet	-	Potential colonization
Eastern Bluebird	Improving	Improving
Veery	Potential extirpation	-
Hermit Thrush	-	Potential colonization
Wood Thrush	Worsening	-
American Robin	Worsening	Improving
Gray Catbird	Worsening*	-
Brown Thrasher	-	Potential colonization
Northern Mockingbird	-	Improving
European Starling	Worsening	Worsening
American Pipit	-	Potential colonization
Cedar Waxwing	Potential extirpation	Improving
Ovenbird	Potential extirpation	-
Prothonotary Warbler	Potential colonization	-
Swainson's Warbler	Potential colonization	-
Kentucky Warbler	Potential colonization	-
Common Yellowthroat	Worsening	-

Common Name	Summer Trend	Winter Trend
Hooded Warbler	Worsening	-
American Redstart	Potential extirpation	-
Northern Parula	Improving	-
Blackburnian Warbler	Potential extirpation	-
Chestnut-sided Warbler	Potential extirpation	-
Palm Warbler	-	Potential colonization [^]
Pine Warbler	-	Potential colonization
Yellow-throated Warbler	Potential colonization	-
Prairie Warbler	Improving	-
Black-throated Green Warbler	Potential extirpation	-
Eastern Towhee	Improving	-
Chipping Sparrow	Worsening	Potential colonization
Field Sparrow	Improving	Improving*
Savannah Sparrow	-	Potential colonization
LeConte's Sparrow	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Song Sparrow	Worsening	Improving
Harris's Sparrow	-	Potential colonization
Dark-eyed Junco	-	Stable
Summer Tanager	Potential colonization	-
Scarlet Tanager	Worsening	-
Northern Cardinal	Improving	Improving
Blue Grosbeak	Potential colonization	-
Indigo Bunting	Improving	-
Dickcissel	Potential colonization	-
Red-winged Blackbird	Stable	-
Brewer's Blackbird	-	Potential colonization
Common Grackle	Worsening	-
Brown-headed Cowbird	Stable	-
Orchard Oriole	Potential colonization	-
Baltimore Oriole	Worsening*	-
House Finch	Worsening*	Worsening*
American Goldfinch	Worsening	Stable
House Sparrow	x	Worsening*