



Little Bighorn Battlefield National Monument

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Little Bighorn Battlefield National Monument (hereafter, the Monument) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Monument, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1).

Among the species likely to be found at the Monument today, climate suitability in summer under the high-emissions pathway is projected to improve for 14, remain stable for 15 (e.g., Figure 2), and worsen for 13 species. Suitable climate ceases to occur for 13 species in summer, potentially resulting in extirpation of those species from the Monument. Climate is projected to become suitable in summer for 19 species not found at the Monument today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 8, remain stable for 4, and worsen for 2 species. Suitable climate ceases to occur for 2 species in winter, potentially resulting in extirpation from the Monument. Climate is projected to become suitable in winter for 49 species not found at the Monument today, potentially resulting in local colonization.

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Monument based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Monument is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.




Figure 1. Projected changes in climate suitability for birds at the Monument, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Monument between the present and 2050 is 0.26 in summer (43rd percentile across all national parks) and 0.31 in winter (48th percentile) under the high-emissions pathway. Potential species turnover declines to 0.17 in summer and 0.22 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Monument is or may become home to 8 species that are highly sensitive to climate change across their range (i.e., they are projected to lose climate suitability in over 50% of their current range in North America in summer and/or winter by 2050; Table 1; Langham et al. 2015).

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Little Bighorn Battlefield National Monument falls within the high turnover group.** Parks anticipating high turnover can focus on actions that increase species' ability to respond to environmental change, such as increasing the amount of potential habitat, working with cooperating agencies and landowners to

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

While the Monument may serve as an important refuge for 7 of these climate-sensitive species, one, the Red Crossbill (*Loxia curvirostra*), might be extirpated from the Monument in summer by 2050.



Figure 2. Climate at the Monument in summer is projected to remain suitable for the Mourning Dove (*Zenaida macroura*) through 2050. Photo by KS Black/Flickr (Public Domain).

improve habitat connectivity for birds across boundaries, managing the disturbance regime, and possibly more intensive management actions. Furthermore, park managers have an opportunity to focus on supporting the 7 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Monument based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Monument is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend
Cackling/Canada Goose	x	Stable
Gadwall	-	Potential colonization
Mallard	Worsening [^]	Improving
Northern Shoveler	-	Potential colonization
Canvasback	-	Potential colonization
Ring-necked Duck	-	Potential colonization
Lesser Scaup	-	Potential colonization
Bufflehead	-	Potential colonization
Hooded Merganser	-	Potential colonization [^]
Scaled Quail	Potential colonization	Potential colonization
Northern Bobwhite	Potential colonization	Potential colonization
Ring-necked Pheasant	Improving	-

Common Name	Summer Trend	Winter Trend
Sharp-tailed Grouse	Worsening [^]	-
Pied-billed Grebe	-	Potential colonization
Eared Grebe	-	Potential colonization
Clark's Grebe	-	Potential colonization
American White Pelican	-	Potential colonization
Great Blue Heron	Potential extirpation	Potential colonization
Black-crowned Night-Heron	-	Potential colonization
Northern Harrier	Worsening [^]	-
Cooper's Hawk	-	Potential colonization
Red-tailed Hawk	Stable	Potential colonization
Ferruginous Hawk	Stable [^]	-
American Coot	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Killdeer	Stable	-
Iceland Gull (Thayer's)	-	Potential colonization
Rock Pigeon	Stable	Potential extirpation
White-winged Dove	-	Potential colonization
Mourning Dove	Stable	-
Greater Roadrunner	-	Potential colonization
Barn Owl	-	Potential colonization
Eastern Screech-Owl	-	Potential colonization
Common Nighthawk	Stable	-
Red-headed Woodpecker	Potential colonization	-
Red-bellied Woodpecker	-	Potential colonization
Ladder-backed Woodpecker	-	Potential colonization
Downy Woodpecker	Improving	Stable
Hairy Woodpecker	Stable	Improving
Northern Flicker	Worsening*	Improving
Gilded Flicker	Potential colonization	-
American Kestrel	x	Improving
Western Wood-Pewee	Worsening*^	-
Say's Phoebe	Worsening	Potential colonization
Ash-throated Flycatcher	Potential colonization	-
Cassin's Kingbird	Improving	-
Western Kingbird	Improving	-
Eastern Kingbird	Stable	-
Loggerhead Shrike	-	Potential colonization
California/Woodhouse's Scrub-Jay (Western Scrub-Jay)	-	Potential colonization
Black-billed Magpie	Worsening^	-

Common Name	Summer Trend	Winter Trend
American Crow	Improving*	Stable
Chihuahuan Raven	Potential colonization	Potential colonization
Common Raven	Potential extirpation	Potential extirpation
Horned Lark	Stable	Improving
Northern Rough-winged Swallow	Improving	-
Tree Swallow	Potential extirpation	-
Barn Swallow	Improving	-
Cliff Swallow	Worsening	-
Black-capped Chickadee	Improving	Worsening*
Bushtit	-	Potential colonization
White-breasted Nuthatch	Improving	Improving
Brown Creeper	-	Potential colonization
Rock Wren	-	Potential colonization
House Wren	Potential extirpation	-
Bewick's Wren	Potential colonization	Potential colonization
Blue-gray Gnatcatcher	Potential colonization	Potential colonization
Ruby-crowned Kinglet	-	Potential colonization
Eastern Bluebird	-	Potential colonization
Western Bluebird	-	Potential colonization
Mountain Bluebird	Worsening	-
American Robin	Stable	Improving
Gray Catbird	Potential extirpation	-
Curve-billed Thrasher	-	Potential colonization
Brown Thrasher	Improving*	-
LeConte's Thrasher	Potential colonization	-

Common Name	Summer Trend	Winter Trend
Crissal Thrasher	Potential colonization	-
Sage Thrasher	-	Potential colonization
Northern Mockingbird	Potential colonization	-
European Starling	Potential extirpation	-
Cedar Waxwing	Stable	Stable
Yellow Warbler	Potential extirpation	-
Yellow-rumped Warbler	-	Potential colonization
Black-throated Gray Warbler	Potential colonization	-
Spotted Towhee	Worsening*	-
Rufous-crowned Sparrow	-	Potential colonization
Canyon Towhee	Potential colonization	Potential colonization
Rufous-winged Sparrow	-	Potential colonization
Cassin's Sparrow	Potential colonization	-
Chipping Sparrow	Potential extirpation	-
Brewer's Sparrow	Worsening*	-
Vesper Sparrow	Potential extirpation	-
Lark Sparrow	Improving	-
Black-throated Sparrow	-	Potential colonization
Lark Bunting	Worsening*	Potential colonization

Common Name	Summer Trend	Winter Trend
Grasshopper Sparrow	Improving*	-
White-crowned Sparrow	-	Potential colonization
Western Tanager	Potential extirpation	-
Blue Grosbeak	Potential colonization	-
Dickcissel	Potential colonization	-
Red-winged Blackbird	Stable	-
Eastern Meadowlark	Potential colonization	-
Western Meadowlark	Stable	-
Brewer's Blackbird	Potential extirpation	-
Common Grackle	Improving*	Potential colonization
Great-tailed Grackle	Potential colonization	-
Brown-headed Cowbird	Stable	Potential colonization
Orchard Oriole	Improving	-
Bullock's Oriole	Worsening	-
Brown-capped Rosy-Finch	-	Potential colonization
House Finch	Stable	Worsening*
Red Crossbill	Potential extirpation [^]	-
Lesser Goldfinch	Potential colonization	-
American Goldfinch	Potential extirpation	Improving