



Lincoln Boyhood National Memorial

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Lincoln Boyhood National Memorial (hereafter, the Memorial) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Memorial, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1). Among the species likely to be found at the Memorial today, climate suitability in summer under the high-emissions pathway is projected to improve for 11, remain stable for 35, and worsen for 14 species. Suitable climate ceases to occur for 19 species in summer, potentially resulting in extirpation of those species from the Memorial (e.g., Figure 2). Climate is projected to become suitable in summer for 19 species not found at the Memorial today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 20, remain stable for 24, and worsen for 6 species. Suitable climate ceases to occur for 5 species in winter, potentially resulting in extirpation from the Memorial. Climate is projected to become suitable in winter for 38 species not found at the Memorial today, potentially resulting in local colonization.

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Memorial based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Memorial is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.




Figure 1. Projected changes in climate suitability for birds at the Memorial, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Memorial between the present and 2050 is 0.20 in summer (32nd percentile across all national parks) and 0.22 in winter (30th percentile) under the high-emissions pathway. Potential species turnover declines to 0.14 in summer and 0.15 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Memorial is or may become home to 2 species that are highly sensitive to climate change across their range (i.e., they are projected to lose climate suitability in over 50% of their current range in North America in summer and/or winter by 2050; Table 1; Langham et al. 2015). While the Memorial may serve as an important refuge for 1 of these

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Lincoln Boyhood National Memorial falls within the intermediate change group.** Parks anticipating intermediate change can best support landscape-scale bird conservation by emphasizing habitat restoration, maintaining natural disturbance regimes, and

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

climate-sensitive species, one, the Mallard (*Anas platyrhynchos*), might be extirpated from the Memorial in summer by 2050.



Figure 2. Although currently found at the Memorial, suitable climate for the American Goldfinch (*Spinus tristis*) may cease to occur here in summer by 2050, potentially resulting in local seasonal extirpation. Photo by John Benson/Flickr (CC BY 2.0).

reducing other stressors. Furthermore, park managers have an opportunity to focus on supporting the 1 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Memorial based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Memorial is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend
Cackling/Canada Goose	x	Stable
Mallard	Potential extirpation [^]	Stable
Blue-winged Teal	-	Potential colonization
Ring-necked Duck	-	Improving
Bufflehead	-	Potential colonization
Northern Bobwhite	Stable	Stable
Wild Turkey	x	Potential extirpation
Eared Grebe	-	Potential colonization
Anhinga	-	Potential colonization
American White Pelican	-	Potential colonization
Great Blue Heron	Stable	-
Great Egret	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Little Blue Heron	Potential colonization	-
Cattle Egret	Potential colonization	-
Green Heron	Stable	-
Yellow-crowned Night-Heron	Potential colonization	-
Black Vulture	Improving	Improving*
Turkey Vulture	x	Improving
Osprey	-	Potential colonization
Mississippi Kite	Improving*	-
Sharp-shinned Hawk	-	Stable
Bald Eagle	-	Stable
Harris's Hawk	Potential colonization	-
Red-shouldered Hawk	Improving	Stable
Red-tailed Hawk	Stable	Improving
Killdeer	Stable	Improving

Common Name	Summer Trend	Winter Trend
Spotted Sandpiper	-	Potential colonization
Greater Yellowlegs	-	Potential colonization
Least Sandpiper	-	Potential colonization
Gull-billed Tern	-	Potential colonization
Forster's Tern	-	Potential colonization
Eurasian Collared-Dove	-	Potential colonization
White-winged Dove	-	Potential colonization
Mourning Dove	Stable	Worsening
Inca Dove	Potential colonization	Potential colonization
Yellow-billed Cuckoo	Improving	-
Greater Roadrunner	Potential colonization	-
Barred Owl	x	Improving
Common Nighthawk	Potential colonization	-
Common Pauraque	-	Potential colonization
Chimney Swift	Worsening	-
Ruby-throated Hummingbird	Stable	-
Belted Kingfisher	Stable	Stable
Red-headed Woodpecker	Stable	Stable
Golden-fronted Woodpecker	Potential colonization	-
Red-bellied Woodpecker	Stable	Improving
Yellow-bellied Sapsucker	-	Improving
Ladder-backed Woodpecker	Potential colonization	-
Downy Woodpecker	Worsening	Worsening
Hairy Woodpecker	Potential extirpation	Worsening
Red-cockaded Woodpecker	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Northern Flicker	Stable	Stable
Pileated Woodpecker	Stable	Stable
American Kestrel	-	Stable
Eastern Wood-Pewee	Worsening	-
Acadian Flycatcher	Stable	-
Willow Flycatcher	Potential extirpation	-
Eastern Phoebe	Stable	-
Great Crested Flycatcher	Stable	-
Brown-crested Flycatcher	Potential colonization	-
Eastern Kingbird	Worsening	-
White-eyed Vireo	Improving	-
Yellow-throated Vireo	Stable	-
Warbling Vireo	Potential extirpation	-
Red-eyed Vireo	Stable	-
Blue Jay	Stable	Stable
American Crow	Stable	Stable
Fish Crow	-	Potential colonization
Northern Rough-winged Swallow	Improving*	-
Purple Martin	Stable	-
Tree Swallow	Potential extirpation	-
Barn Swallow	Stable	-
Cliff Swallow	Potential colonization	-
Carolina Chickadee	Improving*	Improving
Tufted Titmouse	Stable	Improving
Red-breasted Nuthatch	-	Potential extirpation
White-breasted Nuthatch	Worsening*	Worsening*
Brown-headed Nuthatch	Potential colonization^	-
Brown Creeper	-	Worsening

Common Name	Summer Trend	Winter Trend
House Wren	Potential extirpation	Potential colonization
Pacific/Winter Wren	-	Improving
Sedge Wren	-	Potential colonization
Marsh Wren	-	Potential colonization
Carolina Wren	Worsening	Stable
Bewick's Wren	Potential colonization	Potential colonization
Blue-gray Gnatcatcher	Stable	-
Golden-crowned Kinglet	-	Stable
Eastern Bluebird	Improving	Improving
Wood Thrush	Potential extirpation	-
American Robin	Potential extirpation	Stable
Gray Catbird	Potential extirpation	-
Brown Thrasher	Worsening	Improving
Northern Mockingbird	Stable	-
European Starling	Worsening	Stable
American Pipit	-	Potential colonization
Sprague's Pipit	-	Potential colonization
Cedar Waxwing	Potential extirpation	Stable
Chestnut-collared Longspur	-	Potential colonization
Ovenbird	Potential extirpation	-
Worm-eating Warbler	Stable	-
Black-and-white Warbler	Potential colonization	-
Prothonotary Warbler	Stable	-
Swainson's Warbler	Potential colonization	-
Orange-crowned Warbler	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Kentucky Warbler	Improving*	-
Common Yellowthroat	Potential extirpation	Potential colonization
Hooded Warbler	Stable	-
Northern Parula	Stable	-
Yellow Warbler	Potential extirpation	-
Yellow-rumped Warbler	-	Stable
Yellow-throated Warbler	Stable	-
Prairie Warbler	Stable	-
Yellow-breasted Chat	Stable	-
Eastern Towhee	Potential extirpation	x
Bachman's Sparrow	Potential colonization	Potential colonization
American Tree Sparrow	-	Potential extirpation
Chipping Sparrow	Potential extirpation	-
Field Sparrow	Worsening	Improving
Vesper Sparrow	-	Potential colonization
Lark Sparrow	Potential colonization	Potential colonization
Lark Bunting	-	Potential colonization
Fox Sparrow	-	Improving
Song Sparrow	Potential extirpation	Stable
Lincoln's Sparrow	-	Potential colonization
Swamp Sparrow	-	Stable
White-throated Sparrow	-	Improving
Harris's Sparrow	-	Potential colonization
Dark-eyed Junco	-	Stable
Summer Tanager	Stable	-
Scarlet Tanager	Potential extirpation	-
Northern Cardinal	Improving	Improving

Common Name	Summer Trend	Winter Trend
Pyrrhuloxia	-	Potential colonization
Blue Grosbeak	Worsening	-
Indigo Bunting	Stable	-
Painted Bunting	Potential colonization	-
Dickcissel	Stable	-
Red-winged Blackbird	Worsening	Stable
Eastern Meadowlark	Improving	Improving
Western Meadowlark	-	Potential colonization
Brewer's Blackbird	-	Potential colonization
Common Grackle	Worsening	Improving

Common Name	Summer Trend	Winter Trend
Great-tailed Grackle	Potential colonization	Potential colonization
Bronzed Cowbird	-	Potential colonization
Brown-headed Cowbird	Worsening	Improving
Orchard Oriole	Worsening	-
Baltimore Oriole	Stable	-
House Finch	Potential extirpation	Potential extirpation
Purple Finch	-	Potential extirpation
American Goldfinch	Potential extirpation	Stable
House Sparrow	x	Worsening