



Wind Cave National Park

Background

Birds are useful indicators of ecological change because they are highly mobile and generally conspicuous. As climate in a particular place changes, suitability may worsen for some species and improve for others. These changes in climate may create the potential for local extirpation or new colonization. **This brief summarizes projected changes in climate suitability by mid-century for birds at Wind Cave National Park (hereafter, the Park) under two climate change scenarios (see Wu et al. 2018 for full results, and Langham et al. 2015 for more information regarding how climate suitability is characterized).** The high-emissions pathway (RCP8.5) represents a future in which little action is taken to reduce global emissions of greenhouse gases. The low-emissions pathway (RCP2.6) is a best-case scenario of aggressive efforts to reduce emissions. These emissions pathways are globally standardized and established by the Intergovernmental Panel on Climate Change for projecting future climate change. The findings below are model-based projections of how species distributions may change in response to climate change. A 10-km buffer was applied to each park to match the spatial resolution of the species distribution models (10 x 10 km), and climate suitability was taken as the average of all cells encompassed by the park and buffer.

Results

Climate change is expected to alter the bird community at the Park, with greater impacts under the high-emissions pathway than under the low-emissions pathway (Figure 1). Among the species likely to be found at the Park today, climate suitability in summer under the high-emissions pathway is projected to improve for 21, remain stable for 9, and worsen for 16 species. Suitable climate ceases to occur for 42 species in summer, potentially resulting in extirpation of those species from the Park (e.g., Figure 2). Climate is projected to become suitable in summer for 18 species not found at the Park today, potentially resulting in local colonization. Climate suitability in winter under the high-emissions pathway is projected to improve for 8, remain stable for 4, and worsen for 6 species. Suitable climate ceases to occur for 1 species in winter, potentially resulting in extirpation from the Park. Climate is projected to become suitable in winter for 51 species not found at the Park today, potentially resulting in local colonization.

IMPORTANT

This study focuses exclusively on changing climatic conditions for birds over time. But projected changes in climate suitability are not definitive predictions of future species ranges or abundances. Numerous other factors affect where species occur, including habitat quality, food abundance, species adaptability, and the availability of microclimates (see Caveats). Therefore, managers should consider changes in climate suitability alongside these other important influences.

We report trends in climate suitability for all species identified as currently present at the Park based on both NPS Inventory & Monitoring Program data and eBird observation data (2016), plus those species for which climate at the Park is projected to become suitable in the future (Figure 1 & Table 1). This brief provides park-specific projections whereas Wu et al. (2018), which did not incorporate park-specific species data and thus may differ from this brief, provides system-wide comparison and conclusions.




Figure 1. Projected changes in climate suitability for birds at the Park, by emissions pathway and season.

Results (continued)

Potential Turnover Index

Potential bird species turnover for the Park between the present and 2050 is 0.40 in summer (71st percentile across all national parks) and 0.34 in winter (55th percentile) under the high-emissions pathway. Potential species turnover declines to 0.27 in summer and 0.23 in winter under the low-emissions pathway. Turnover index was calculated based on the theoretical proportions of potential extirpations and potential colonizations by 2050 relative to today (as reported in Wu et al. 2018), and therefore assumes that all potential extirpations and colonizations are realized. According to this index, no change would be represented as 0, whereas a complete change in the bird community would be represented as 1.

Climate Sensitive Species

The Park is or may become home to 14 species that are highly sensitive to climate change across their range (i.e., they are projected to lose climate suitability in over 50% of their current range in North America in summer and/or winter by 2050; Table 1; Langham et al. 2015). While the Park may serve as an important refuge for 6 of these

Management Implications

Parks differ in potential colonization and extirpation rates, and therefore different climate change adaptation strategies may apply. **Under the high-emissions pathway, Wind Cave National Park falls within the high turnover group.** Parks anticipating high turnover can focus on actions that increase species' ability to respond to environmental change, such as increasing the amount of potential habitat, working with cooperating agencies and landowners to improve habitat connectivity

Caveats

The species distribution models included in this study are based solely on climate variables (i.e., a combination of annual and seasonal measures of temperature and precipitation), which means there are limits on their interpretation. Significant changes in climate suitability, as measured here, will not always result in a species response, and all projections should be interpreted as potential trends. Multiple other factors mediate responses to climate change, including habitat availability, ecological processes

climate-sensitive species, 8 might be extirpated from the Park in at least one season by 2050.



Figure 2. Although currently found at the Park, suitable climate for the American Robin (*Turdus migratorius*) may cease to occur here in summer by 2050, potentially resulting in local seasonal extirpation. Photo by Andy Reago & Chrissy McClarren/Flickr (CC BY 2.0).

for birds across boundaries, managing the disturbance regime, and possibly more intensive management actions. Furthermore, park managers have an opportunity to focus on supporting the 6 species that are highly sensitive to climate change across their range (Table 1; Langham et al. 2015) but for which the park is a potential refuge. Monitoring to identify changes in bird communities will inform the selection of appropriate management responses.

that affect demography, biotic interactions that inhibit and facilitate species' colonization or extirpation, dispersal capacity, species' evolutionary adaptive capacity, and phenotypic plasticity (e.g., behavioral adjustments). Ultimately, models can tell us where to focus our concern and which species are most likely to be affected, but monitoring is the only way to validate these projections and should inform any on-the-ground conservation action.

More Information

For more information, including details on the methods, please see the scientific publication ([Wu et al. 2018](#)) and the [project overview brief](#), and visit the [NPS Climate Change Response Program website](#).

References

eBird Basic Dataset (2016) Version: ebd_relAug-2016. Cornell Lab of Ornithology, Ithaca, New York.

Langham et al. (2015) Conservation Status of North American Birds in the Face of Future Climate Change. PLOS ONE.

Wu et al. (2018) Projected avifaunal responses to climate change across the U.S. National Park System. PLOS ONE.

Contacts

Gregor Schuurman, Ph.D.
Ecologist, NPS Climate Change Response Program
970-267-7211, gregor_schuurman@nps.gov

Joanna Wu
Biologist, National Audubon Society
415-644-4610, science@audubon.org

Species Projections

Table 1. Climate suitability projections by 2050 under the high-emissions pathway for all birds currently present at the Park based on both NPS Inventory & Monitoring Program data and eBird observation data, plus those species for which climate at the Park is projected to become suitable in the future. "Potential colonization" indicates that climate is projected to become suitable for the species, whereas "potential extirpation" indicates that climate is suitable today but projected to become unsuitable. Omitted species were either not modeled due to data deficiency or were absent from the I&M and eBird datasets. Observations of late-season migrants may result in these species appearing as present in the park when they may only migrate through. Species are ordered according to taxonomic groups, denoted by alternating background shading.

* Species in top and bottom 10th percentile of absolute change

^ Species that are highly climate sensitive

- Species not found or found only occasionally, and not projected to colonize by 2050

x Species not modeled in this season

Common Name	Summer Trend	Winter Trend	Common Name	Summer Trend	Winter Trend
Wood Duck	-	Potential colonization		colonization	colonization
Gadwall	-	Potential colonization	Sharp-tailed Grouse	Potential extirpation [^]	-
Mallard	Worsening* [^]	Improving	Wild Turkey	x	Stable
Blue-winged Teal	Worsening	-	Pied-billed Grebe	-	Potential colonization
Northern Shoveler	-	Potential colonization	Eared Grebe	-	Potential colonization
Canvasback	-	Potential colonization	Western Grebe	-	Potential colonization
Ring-necked Duck	-	Potential colonization	American White Pelican	-	Potential colonization
Lesser Scaup	-	Potential colonization	Great Blue Heron	Stable	Potential colonization
Bufflehead	-	Potential colonization	Black-crowned Night-Heron	-	Potential colonization
Hooded Merganser	-	Potential colonization [^]	Northern Harrier	Worsening [^]	-
Scaled Quail	Potential colonization	Potential colonization	Cooper's Hawk	x	Potential colonization
Gambel's Quail	-	Potential colonization	Bald Eagle	-	Worsening*
Northern Bobwhite	Potential	Potential	Red-tailed Hawk	Worsening	-
			Rough-legged Hawk	-	Worsening*

Common Name	Summer Trend	Winter Trend
American Coot	-	Potential colonization
Killdeer	Improving	-
Upland Sandpiper	Worsening*	-
Long-billed Curlew	Worsening^	-
Ring-billed Gull	Potential extirpation^	-
Rock Pigeon	Potential extirpation	-
Eurasian Collared-Dove	-	Potential colonization
Mourning Dove	Improving	Improving
Greater Roadrunner	-	Potential colonization
Barn Owl	-	Potential colonization
Great Horned Owl	x	Stable
Burrowing Owl	Improving*^	-
Common Nighthawk	Improving	-
Black-chinned Hummingbird	Potential colonization	-
Belted Kingfisher	Potential extirpation	-
Red-headed Woodpecker	Improving*	-
Red-bellied Woodpecker	-	Potential colonization
Red-naped Sapsucker	Potential extirpation^	-
Ladder-backed Woodpecker	-	Potential colonization
Downy Woodpecker	Improving	-
Hairy Woodpecker	Potential extirpation	Worsening*
Northern Flicker	Potential extirpation	Improving
Gilded Flicker	Potential colonization	-
Western Wood-Pewee	Potential extirpation^	-
Willow Flycatcher	Potential extirpation	-

Common Name	Summer Trend	Winter Trend
Least Flycatcher	Potential extirpation	-
Dusky Flycatcher	Worsening	-
Cordilleran Flycatcher	Worsening	-
Eastern Phoebe	Improving	-
Say's Phoebe	Worsening	Potential colonization
Cassin's Kingbird	Potential colonization	-
Western Kingbird	Improving	-
Eastern Kingbird	Worsening*	-
Scissor-tailed Flycatcher	Potential colonization	-
Loggerhead Shrike	-	Potential colonization
Bell's Vireo	Improving	-
Warbling Vireo	Improving	-
Red-eyed Vireo	Potential extirpation	-
Gray Jay	Potential extirpation	-
Pinyon Jay	Stable	-
Blue Jay	Improving*	-
Black-billed Magpie	Worsening^	Worsening*
Clark's Nutcracker	Potential extirpation^	-
American Crow	Stable	Improving
Chihuahuan Raven	Potential colonization	Potential colonization
Horned Lark	Stable	-
Northern Rough-winged Swallow	Improving	-
Tree Swallow	Potential extirpation	-
Violet-green Swallow	Potential extirpation	-
Barn Swallow	Improving	-
Cliff Swallow	Worsening	-
Black-capped Chickadee	Stable	Worsening*
Juniper Titmouse	Potential	-

Common Name	Summer Trend	Winter Trend
	colonization	
Bushtit	-	Potential colonization
Red-breasted Nuthatch	Potential extirpation	Potential extirpation
White-breasted Nuthatch	Stable	Stable
Pygmy Nuthatch	Potential colonization	-
Brown Creeper	Potential extirpation^	Potential colonization
Rock Wren	Worsening	Potential colonization
House Wren	Worsening*	-
Marsh Wren	-	Potential colonization
Bewick's Wren	Potential colonization	Potential colonization
Ruby-crowned Kinglet	-	Potential colonization
Eastern Bluebird	Improving	Potential colonization
Mountain Bluebird	Potential extirpation	-
Townsend's Solitaire	Potential extirpation^	Worsening*
Swainson's Thrush	Potential extirpation	-
American Robin	Potential extirpation	Improving
Curve-billed Thrasher	-	Potential colonization
Brown Thrasher	Improving	Potential colonization
Sage Thrasher	-	Potential colonization
Northern Mockingbird	Potential colonization	-
European Starling	Potential extirpation	-
Cedar Waxwing	Potential extirpation	Stable
Chestnut-collared Longspur	-	Potential colonization

Common Name	Summer Trend	Winter Trend
Smith's Longspur	-	Potential colonization
Ovenbird	Potential extirpation	-
Common Yellowthroat	Potential extirpation	-
American Redstart	Potential extirpation	-
Yellow Warbler	Potential extirpation	-
Yellow-rumped Warbler	Potential extirpation	Potential colonization
Yellow-breasted Chat	Potential extirpation	-
Spotted Towhee	Potential extirpation	-
Rufous-crowned Sparrow	-	Potential colonization
Rufous-winged Sparrow	Potential colonization	Potential colonization
Cassin's Sparrow	Potential colonization	-
American Tree Sparrow	-	Improving
Chipping Sparrow	Potential extirpation	-
Brewer's Sparrow	Potential extirpation	-
Field Sparrow	Potential extirpation	-
Vesper Sparrow	Potential extirpation	-
Lark Sparrow	Improving	-
Lark Bunting	Worsening	-
Grasshopper Sparrow	Improving	-
Song Sparrow	Potential extirpation	Potential colonization
Lincoln's Sparrow	-	Potential colonization
White-crowned Sparrow	-	Potential colonization
Dark-eyed Junco	x	Improving
Western Tanager	Stable	-

Common Name	Summer Trend	Winter Trend
Northern Cardinal	Improving	Potential colonization
Black-headed Grosbeak	Potential extirpation	-
Blue Grosbeak	Potential colonization	-
Lazuli Bunting	Potential extirpation	-
Indigo Bunting	Improving	-
Dickcissel	Potential colonization	-
Red-winged Blackbird	Worsening	Potential colonization
Eastern Meadowlark	Potential colonization	-
Western Meadowlark	Stable	-
Rusty Blackbird	-	Potential colonization
Brewer's Blackbird	Potential	-

Common Name	Summer Trend	Winter Trend
	extirpation	
Common Grackle	Improving	Potential colonization
Great-tailed Grackle	Potential colonization	Potential colonization
Brown-headed Cowbird	Potential extirpation	Potential colonization
Orchard Oriole	Improving*	-
Bullock's Oriole	Stable	-
Baltimore Oriole	Potential colonization	-
Red Crossbill	Potential extirpation [^]	x
Pine Siskin	Potential extirpation	-
American Goldfinch	Potential extirpation	Improving