



2015 Voyageurs National Park Moose Population Survey Report

Natural Resource Data Series NPS/VOYA/NRDS—2015/XXX



ON THE COVER

Bull moose with deformed antlers observed during the 2015 aerial survey in Voyageurs National Park. The same moose was seen in 2014, based on similar antler development. The cause of the deformity is unknown but similar deformities have been linked to heavy loads of carotid (or arterial) worms in other moose populations.

2015 Voyageurs National Park Moose Population Survey Report

Natural Resource Data Series NPS/VOYA/NRDS—2015/XXX

Steve K. Windels and Bryce T. Olson

National Park Service
Voyageurs National Park
360 Hwy 11 East
International Falls, MN 56649

April 2015

U.S. Department of the Interior
National Park Service
Natural Resource Stewardship and Science
Fort Collins, Colorado

The National Park Service, Natural Resource Stewardship and Science office in Fort Collins, Colorado, publishes a range of reports that address natural resource topics. These reports are of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Data Series is intended for the timely release of basic data sets and data summaries. Care has been taken to assure accuracy of raw data values, but a thorough analysis and interpretation of the data has not been completed. Consequently, the initial analyses of data in this report are provisional and subject to change.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner.

This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U.S. Government.

This report is available from the Natural Resource Publications Management website (<http://www.nature.nps.gov/publications/nrpm/>).

Please cite this publication as:

Windels, S. K., and B.T. Olson. 2015. 2015 Voyageurs National Park Moose Population Survey Report. Natural Resource Data Series NPS/VOYA/NRDS—2015/XXX. National Park Service, Fort Collins, Colorado.

Contents

	Page
Contents.....	iii
Figures.....	iii
Tables.....	iii
Acknowledgments.....	iv
Introduction.....	5
Methods.....	6
Results.....	8
Discussion.....	10
Literature Cited.....	12

Figures

	Page
Figure 1. 2015 moose survey area in Voyageurs National Park, Minnesota, USA.....	4

Tables

	Page
Table 1. Population estimates and demographic characteristics for moose in the Kabetogama Peninsula, Voyageurs National Park, USA, derived from aerial surveys and other sources from 1991-2015.....	5

Acknowledgments

Funding for this project was provided by Voyageurs National Park. We also acknowledge support from the University of Minnesota (Duluth and Twin Cities campuses) for GPS collars and data acquisition for collared moose used in the visibility trials. Ron Moen, University of Minnesota-Duluth, is a co-investigator for the ongoing moose research in VOYA, including the capture and collaring of moose in the park since 2010. Special thanks to Voyageurs National Park Pilot Steve Mazur for assistance with data collection for the 2015 survey. We also appreciate the support of the Minnesota Department of Natural Resources Wildlife Health Program and the University of Minnesota Veterinary Diagnostic Lab for providing pregnancy testing and pathological examinations of dead moose. Ron Moen, Mike Schrage, and Mary Graves provided helpful comments on an earlier draft of this report.

Introduction

Voyageurs National Park (Minnesota) was established in 1975 in part to fulfill the National Park Service's mission to preserve and protect wildlife populations and provide opportunities for the public to enjoy them. Moose (*Alces alces*) are native to Voyageurs National Park (VOYA) but recent declines in other moose populations in the region raised concerns about the long-term viability of moose in the park. Moose populations in northwestern Minnesota declined precipitously during the period 1984-2000 (Murray et al. 2006). Moose populations in northeastern Minnesota have been experiencing similar declines in recent years, with recent estimates nearly 50% lower than the 2006 estimate (DelGiudice 2015). Voyageurs National Park is not surveyed as part of the state's systematic annual survey because it lies just outside of primary moose range in northeastern Minnesota (Fig. 1; DelGiudice 2015). Voyageurs National Park, in collaboration with the University of Minnesota-Duluth, began more intensive monitoring and research of moose in and adjacent to the park in 2009 to better understand local moose population dynamics. Information from these efforts will help NPS managers ensure the survival of moose in VOYA for future generations.

Methods

We surveyed the moose population within the boundaries of Voyageurs National Park during Feb. 2-14, 2015. The survey area was limited to the Kabetogama Peninsula, a 305 km² roadless area in the center of the park where >90% of the Park's moose population occurs (Figure 1). Surveys were conducted using a 2-seat Aviat Husky during which the pilot and observer search for moose while the plane flies in overlapping circles at an intensity of at least 3.5 min/km². The peninsula was broken down into 23 separate survey units to facilitate the completion of the survey, and all units were surveyed. For each observed moose we recorded location, group size, sex/age class (calf, yearling, adult cow, adult bull, unknown), and whether the animal was standing or bedded. We also recorded all observations of white-tailed deer (*Odocoileus virginianus*) and gray wolves (*Canis lupus*).

We conducted 18 test plots to estimate visibility (detection probability) of moose for our survey method. We searched test plots, using the same flight pattern and intensity as the survey plots, for moose wearing GPS telemetry collars. Locations of moose not observed during the test plots were confirmed by GPS locations or using VHF telemetry. Test plots were completed during Feb. 1-Mar. 11, 2015 during conditions similar to those that occurred during the survey.

The number of moose observed during the aerial survey was adjusted with the estimated detection probability, giving a population estimate for the Kabetogama Peninsula (\pm 90% Confidence Intervals) during the survey period. We also estimated other measures of population status, including calf:cow ratio, twinning rates, and bull:cow ratio. No moose were captured in 2015 and therefore we did not estimate pregnancy rates (percent of adult females that were pregnant via blood progesterone levels) as was done in previous years.




Figure 1. 2015 moose survey area in Voyageurs National Park, Minnesota, USA. The Kabetogama Peninsula (305 km²) contains >95% of the Park's moose population. A small pocket of moose (approx. 10-15 individuals) also exists west of the park in the Rat Root Lake area, and evidence from GPS collars suggests that some moose seasonally move between this area and the Kabetogama Peninsula. Some moose also reside in the southeastern portion of the park but we did not survey this part of the park in 2015.

Results

Survey conditions were considered “good” to “excellent” during the 2015 survey (including visibility trials), with snow depths exceeding 35 cm (14”) throughout the Kabetogama Peninsula and little snow in the canopies of trees. We detected 76% ($\pm 9\%$; 16 out of 21) of collared moose during visibility trials.

We counted 34 moose during the survey (13 bulls, 13 cows, 3 unknown adults, 1 yearling, and 4 calves). After correcting for visibility, the 2015 population estimate for the Kabetogama Peninsula was 46 moose (90% Confidence Interval = 41-52), or 0.15 moose/km² (0.39 moose/mi²). We accounted for one additional moose (1 calf) not observed during the survey but later observed with its collared mother on the Kabetogama Peninsula. The minimum number of moose on the Kabetogama Peninsula during the 2015 survey based on known individuals was 35, close to the lower end of the 90% Confidence Interval for the survey estimate. Additionally, 1 collared bull and 1 collared cow left the Kabetogama Peninsula for adjacent Ontario in early winter before the survey began. The 2015 population estimate is similar to those from the period 2009-2014 (Table 1). Indices of calf production in 2015 were relatively low, similar to 2014. No twins were observed in 2015, as has been the case in most survey years. The estimated calf:cow ratio was 0.38 and calves were 14% of the population. The bull:cow ratio observed during the 2015 survey (1.0) was higher than in previous years, but is likely a reflection of our sampling protocol rather than an actual change in the adult sex ratio over the last year.

Table 1. Population estimates and demographic characteristics for moose in the Kabetogama Peninsula, Voyageurs National Park, USA, derived from aerial surveys and other sources from 1991-2015.

Year	Population Estimate	90% Confidence Interval for Estimate	Calves: Cow	% Calves	% Twins ^a	Bulls: Cow	% Pregnant ^b
1991	31	23-57	-	9	-	-	-
1992	47	35-72	-	9	-	-	-
1997	53	32-88	-	25	~10 ^c	-	-
1998	38	23-63	-	9	0	-	-
2009	51	44-58	-	7	0	-	-
2010	41	36-47	0.54	23	0	0.82	60
2011	45	39-51	0.60	28	8	0.53	69
2012	-----Not Surveyed-----						33
2013	46	43-50	0.61	25	6	0.56	63
2014	40	34-48	0.23	11	0	0.46	-
2015	46	41-52	0.38	14	0	1.00	-

^aPercentage of twins observed among all cows.

^bEstimated from serum progesterone levels from blood samples collected during winter capture for GPS collaring.

^cOne set of twins recorded; % Twins for 1997 based on assumption of 1:1 adult sex ratio.

Discussion

Voyageurs National Park staff have been monitoring the park's moose population since 2009, including conducting aerial surveys to estimate population size and demographics and monitoring adult moose with GPS collars to understand survival, habitat use, and other behaviors. Aerial survey data support that VOYA maintains a stable, low density moose population in the Kabetogama Peninsula. Since Feb. 2010, we have monitored adult survival with GPS/VHF telemetry collars on 25-50% of the adult population. All 10 collared moose survived the period Mar. 1, 2014 to Feb. 28, 2015. Three additional collared moose survived at least 2-10 months until we lost contact with their collars. Overall, we have confirmed death on 7 of 22 adult moose collared since 2010, resulting in a mean annual mortality rate of approximately 7% during the period Feb. 1, 2010-Jan. 31, 2015 (range = 0-16% per year; Voyageurs National Park, unpublished data). Our estimate of annual adult mortality is similar to those reported for non-hunting mortality rates from other moose populations in North America (Van Ballenberghe and Ballard 2007) but noticeably less than those reported for the northwestern Minnesota population in 1995-2000 (21%; Murray et al. 2006) and for the northeastern Minnesota population in 2002-2008 (19%; Lenarz et al. 2009) and 2013-2014 (11-20%, M. Carstensen, personal communication). Survey data from 2009-2015 suggest that reproduction and recruitment may be enough to offset observed adult mortality, and therefore maintain a stable population (Windels 2014).

A recent study suggested that increased wolf abundance and hence increased predation may be partially responsible for the dramatic decline in the northeastern moose population since 2006 (Mech and Fieberg 2014). There is little evidence to support that wolves are having a similar negative impact on the moose population in VOYA at present despite a relatively high wolf:moose ratio (16 wolves : 46 moose; VOYA, unpublished data) on the Kabetogama Peninsula. As stated above, adult survival in VOYA is very high and the population has remained stable for the last 7 years. Causes of death of the 6 non-capture related cases for adult collared moose have been health-related (50%) or unknown (50%), though predation could be responsible for some of the cases where scavengers had consumed too much of the carcass to determine the original cause of death.

Wolves and black bears, both of which are abundant in Voyageurs, readily prey on young calves and annual survival of calves in most moose populations is low when bears and wolves are present. Productivity data suggest that a relatively low proportion of cow moose have calves in any given year (4-year mean = 62%, Table 1) but those calves that are born have a reasonable chance of survival (4-year mean \approx 40-45%) to their first birthday (VOYA, unpublished data). Our estimate of 1-year survival is similar to that reported for 2002-2008 in northeastern Minnesota (Lenarz et al. 2010) and slightly higher than reported in a current study in the same area (Severud et al. 2014). A recent study of wolf diet in Minnesota based on scat analysis confirms that wolves in VOYA rarely consume moose relative to other areas of northeastern MN; 5-13% of biomass of VOYA wolf diets in summer and winter were comprised of moose versus 32-54% in other areas. Wolves in VOYA consume primarily deer (53-78% of biomass) and beaver (7-26%) (Ibrahim et al., unpublished data). Densities of deer (\sim 10 deer/mi²) and beavers (\sim 13 beavers/mi²) are relatively high in the VOYA area (VOYA, unpublished data) and wolves and other potential predators typically prefer to prey on deer and beavers rather than

moose. If abundance of alternative prey such as deer and beavers dramatically drop, however, we might expect that predation on adult or calf moose may increase.

In addition to population monitoring, Voyageurs National Park is currently investigating other aspects of moose ecology in collaboration with the University of Minnesota-Duluth, Minnesota Zoo, Bemidji State University, Lakehead University, University of Minnesota-Twin Cities, and other partners. Recently completed projects in the park include a study to understand factors that affect temperature in different types of moose habitat (Olson et al. 2014); a study of spatial patterns in deer infected with brainworm, a parasite fatal to moose (VanderWaal et al. 2015); and a study of abundance of terrestrial gastropods, the main vector of brainworm for deer and moose, in different forest habitats (Cyr et al. 2015). Ongoing studies include understanding how moose behave in response to high temperatures and other weather events; identifying potential thermal refugia for moose in the face of a changing climate; interactions of moose, deer, beavers, and wolves; and how and why moose use wetlands for foraging and temperature regulation.

Literature Cited

- Cyr, T., S.K. Windels, R.A. Moen, and J.W. Warmbold. 2014. Diversity and abundance of terrestrial gastropods in Voyageurs National Park, MN: implications for risk of moose to *Parelaphostrongylus tenuis* infection. *Alces* 50: 121-132.
- DelGiudice, G.D. 2015. Aerial Moose Survey. Minnesota Department of Natural Resources. http://files.dnr.state.mn.us/wildlife/moose/2015_mooseurvey.pdf (accessed April 14, 2015).
- Lenarz, M. S., M.E. Nelson, M.W. Schrage, and A.J. Edwards. 2009. Temperature mediated moose survival in northeastern Minnesota. *Journal of Wildlife Management* 73: 503-510.
- Lenarz, M. S., Fieberg, J., Schrage, M. W., & Edwards, A. J. (2010). Living on the edge: viability of moose in northeastern Minnesota. *The Journal of Wildlife Management* 74: 1013-1023.
- Mech, L.D., and J. Fieberg. 2014. Re-evaluating the northeastern Minnesota moose decline and the role of wolves. *Journal of Wildlife Management* 78:1143-1150.
- Murray, D.J., E.W. Cox, W.B. Ballard, H.A. Whitlaw, M.S. Lenarz, T.W. Custer, T. Barnett, and T.K. Fuller. 2006. Pathogens, nutritional deficiency, and climate influences on a declining moose population. *Wildlife Monographs* 166:1-30.
- Olson, B.T., S.K. Windels, M. Fulton, and R.A. Moen. 2014. Fine-scale temperature patterns in the southern boreal forest: implications for the cold-adapted moose. *Alces* 50: 105-121.
- Severud, W. J., G. D. DelGiudice, T. R. Obermoller, K. J. Foshay, and R. G. Wright. 2014. Using GPS collars to determine moose calving and cause-specific mortality of calves in northeastern Minnesota: progress report on second field season. Pages 40-56 in L. Cornicelli, M. Carstensen, M. D. Grund, M. A. Larson, and J. S. Lawrence, editors. *Summaries of wildlife research findings, 2013*. Minnesota Department of Natural Resources, St. Paul.
- Van Ballenberghe, V., and W.B. Ballard. 2007. Population dynamics. Pages 223-245 in A.W. Franzmann and C.C. Schwartz, editors. *Ecology and Management of the North American Moose*, 2nd ed. University Press of Colorado, Boulder, CO.
- VanderWaal, K.L., S.K. Windels, B.T. Olson, T. Vannatta, and R.A. Moen. 2014. Spatial epidemiology of liver fluke and meningeal worm in white-tailed deer in northern Minnesota, USA. *Parasitology*: 1-13.
- Windels, S. K. 2014. 2014 Voyageurs National Park moose population survey report. Natural Resource Data Series NPS/VOYA/NRDS—2014/645. National Park Service, Fort Collins, Colorado.